

The millipede genus *Strongylosoma* in the Caucasus (Diplopoda: Polydesmida, Paradoxosomatidae)

Aleksandr EVSYUKOV¹⁾, Sergei GOLOVATCH²⁾ & Hans S. REIP³⁾

¹⁾ Lyceum No. 1 “Classical”, Balakireva Str. 32, Rostov-on-Don 344004, Russia;
e-mail: aevsukov@mail.ru

²⁾ Institute for Problems of Ecology and Evolution, Russian Academy of Science, Leninsky pr. 33,
Moscow 119071, Russia; e-mail: sgolovatch@yandex.ru

³⁾ Senckenberg Museum of Natural History Görlitz, Am Museum 1,
D-02826 Görlitz, Germany; e-mail: reip@myriapoden-info.de
Corresponding author: sgolovatch@yandex.ru

Received 19 August 2014; accepted 6 January 2015
Published 12 April 2016

Abstract. Only two species of *Strongylosoma* Brandt, 1833 inhabit the Caucasus: *Strongylosoma kordylamythrum* Attems, 1898 and *S. lenkoranum* Attems, 1898. Both are quite distinct morphologically, their main diagnostic features are in the structure of the gonopod telopodite and shape of the ventral hump on male femur 3. The former species occurs in most of the Caucasus except for the region’s eastern parts where only the latter species occurs. Parapatry is recorded only in Mountainous Karabakh and Hyrcania (within both southeastern Azerbaijan and northwestern Iran), but even there these species are strictly allopatric, in never being found together. Their distributions are refined and mapped, based both on records in the literature and an abundance of new samples.

Key words. Distribution, map, diagnostic features, Diplopoda, *Strongylosoma kordylamythrum*, *Strongylosoma lenkoranum*, Caucasus.

INTRODUCTION

The rather large, eastern Euro-Mediterranean millipede genus *Strongylosoma* Brandt, 1833 contains 13 unquestioned species or subspecies, which are confined to Central, Eastern and Southeastern Europe, the Caucasus and the Middle East. A further 30 species, largely tropical, are dubious (Nguyen & Sierwald 2013). For a long time only two species have been recorded as occurring in the Caucasus: *Strongylosoma kordylamythrum* Attems, 1898 and *S. lenkoranum* Attems, 1898 (e.g. Attems 1898, Lohmander 1936), but their distributions have hitherto remained rather obscure. The ubiquitous, definitely introduced *Oxidus gracilis* (C. L. Koch, 1847) is the only other member of the family Paradoxosomatidae recorded in the Caucasus (e.g. Lignau 1915).

The present paper is based on numerous new faunistic records, and summarizes and updates the nomenclatural history and distributions of both *Strongylosoma* species in the Caucasus.

MATERIAL AND METHODS

A large quantity of material of both *S. kordylamythrum* and *S. lenkoranum* was amassed by the second author, which is now mostly housed in the Zoological Museum of Moscow State University (ZMUM), Moscow, Russia. Additional samples were accumulated by the first and third authors, which are now in their private collections in Rostov-on-Don and Jena, referred to below as (AE) and (CHR), respectively. A further few samples are in the Staatliches Museum für Naturkunde, Görlitz (SMNG) and one in the Natural History Museum of Denmark, Copenhagen (ZMUC). This material allows not only serious refinements of the distributions of both these species, but also a clearer morphological diagnosis of both these species.

The following abbreviations are used: D – description or descriptive notes, R – new records, M – mere mention, ind. – individual, juv. – juvenile form.

TAXONOMIC PART

Strongylosoma kordylamythrum Attems, 1898 (Figs 1, 2, 5)

Strongylosoma kordylamythrum Attems, 1898: 312 (D).

Strongylosoma kordylamythrum: Lignau 1903: 32 (D, R), 1915: 376 (D, R); Attems 1914: 227 (M), 1926: 247 (D), 1937: 38, 39 (D); Verhoeff 1921: 40, 42 (D, R), 1940: 37 (D); Lohmander 1936: 32 (D, R); Lang 1959: 1795 (R), 1964: 241 (R); Kobakhidze 1965: 391 (R); Jeekel 1968: 94 (M); Ljovuschkin & Lokshina, 1975: 211 (R); Golovatch 1983: 167 (R); Talikadze 1984: 143 (M); Bababekova 1996: 90 (R); Enghoff & Moravvej 2005: 68 (R); Golovatch & Matyukhin 2011: 115 (R); Evsyukov & Golovatch 2013: 209 (D, R); Nguyen & Sierwald 2013: 1268 (M); Zuev 2014: 348 (R).

Strongylosoma cordylamythrum (lapsus calami!): Muralewicz 1907: 340 (M, R), 1911: 10 (M), 1913: 220 (R); Ghilarov 1972: 38 (R).

Strongylosoma pallipes (misidentification): Muralewicz 1927: 7 (M).

Strongylosoma lenkoranum (misidentification): Lohmander 1932: 3 (M).

MATERIAL EXAMINED. **Russia:** 2 ♂♂, 1 ♀ (ZMUM p2480). **Chechenia:** Argun River valley, 5 km N of Shatoi, *Corylus*, *Fagus*, *Carpinus* etc. forest, 750 m a. s. l., litter, under stones & bark, 18 July 1986; 2 ♂♂, 1 ♀ (ZMUM p2481). **Ingushetia:** Barsuki near Nazran, *Crataegus* & *Fraxinus* grove along road, litter & under stones, 6 June 1982; 1 ♀ (ZMUM p2482), Assa River valley, ca. 9 km SSW of Muzhichi, 800 m a. s. l., *Fagus*, *Alnus*, *Carpinus* etc. forest, litter, under bark & stones, 15 July 1986, all leg. S. Golovatch; 1 ♂ (AE). **Dagestan:** Makhachkala, park, 1 December 2012, leg. Galimova; 1 ♀ (AE), Makhachkala, 14 April 2012; 1 ♀ (AE), Khindakh, 11 June 2012, all leg. E. Il'yna; 1 ♂, 1 ♀ (ZMUM p2483). **Krasnodar Prov.:** Sochi, Khosta, *Taxus* & *Buxus* grove with *Fagus*, litter & under bark, 15 May 1985; 1 ♀ (ZMUM p2484), Caucasian Nature Reserve, Pslukh, 20 km E of Krasnaya Polyana, Mt Kogot, *Fagus* & *Abies* forest, 1400 m a. s. l., litter, under bark & stones, 18–20 May 1985; 3 ♂♂ (ZMUM p2485), Caucasian Nature Reserve, Krasnaya Polyana, 600–750 m a. s. l., *Quercus*, *Fagus*, *Castanea*, *Caprinus* etc. forest, litter, bark, stones, 8–9 August 1986; 1 ♂, 1 ♀, (ZMUM p2486), Sochi, Khosta, *Taxus* & *Buxus* grove, litter, 28 October 1981; 4 ♀♀ (ZMUM p2487), Tuapse Distr., 15 km SE of Novomikhailovskiy, Psebe, deciduous forest, under stones, litter & in rotten logs, 29 October 1981; 1 ♀, 1 juv. (ZMUM p2488), same locality, Mt Fatse, 28 August 1974, all leg. S. Golovatch; 5 ♀♀ (ZMUM p2489), near Psebe, cave, 11 July 1961, leg. S. Ljovuschkin; 1 ♂, 2 ♀♀, 1 juv. (ZMUM p2490), Sochi, ca. 8 km E of Khosta, Cave “Our Lady”, 16 May 1985; 2 ♀♀ (ZMUM p2491), same locality, *Buxus*, *Fagus*, *Acer* etc. forest, litter, under stones, 18 May 1985; 3 ♂♂, 2 ♀♀ (ZMUM p2492), Severskaya Distr., 2–10 km S of Ubinskaya, *Quercus*, *Fagus*, *Carpinus* etc. forest, 300–450 m a. s. l., litter & under bark, 3 & 4 July 1986, all leg. S. Golovatch; 1 ♀, 1 juv. (ZMUM p2493), near Khosta, Cave Labirintovaya, 27 November 1985, leg. N. Myuge; 1 ♂ (ZMUM p2494), Krasnaya Polyana, 6–8 April 1978, leg. V. Dolin; 10 ♂♂, 4 ♀♀ (ZMUM p2495). **Adygea:** Caucasian Nature Reserve, Pasture Abago near Guzeripl, *Abies*, *Fagus*, *Acer*, *Betula* etc. forest, up to timber line & in subalpine meadow, 1700–1850 m a. s. l., litter, under bark & stones, 24–26 May 1985; 1 ♂, 1 ♀ (ZMUM p2496), same locality, *Abies* & *Fagus* forest, 1350–1400 m a. s. l., litter, under bark & stones, 24–26 May 1985, all leg. S. Golovatch; 1 ♀ (AE), near Kamennomostskiy, Meshokho River valley, under stones, 26 June 2011, leg. D. Khisametdinova; 1 ♀ (AE), same locality, litter, 17 June 2007, leg. A. Evsyukov; 1 ♀ (AE), near Nikel, Belaya River valley, 28 June 2007, leg. G. Chesnokov; 1 ♂ (AE), same locality, 18 June 2010; 1 ♀ (AE), same locality, under logs, 28 August 2009, all leg. A. Evsyukov; 1 ♂ (AE), same locality, 18 June 2010, leg. I. Gorbenko; 2 ♂♂, 2 ♀♀ (AE), Caucasian Nature Reserve, Lagonaki (= Lago-Naki) Plateau, 24 June 2010, leg. A. Evsyukov & D. Khisametdinova; 1 ♀ (AE), Maikop, 14 March 2008, leg. M. Narbtev; 2 ♀♀ (AE), near Kamennomostskiy, 16 June 2012, leg. M. Shapovalov; 1 ♂ (AE), Azish-Tau Mt Ridge, forest, under stones, 10 May 2013, leg. Y. Kochetov & D. Khisametdinova; 1 ♂, 1 ♀ (CHR), Lagonaki (= Lago-Naki) Plateau, source area of Kurdships River, 44° 04' 45.1" N, 40° 00' 00" E, 16 August 2012; 1 ♀ (CHR), Urushrent River valley, right bank, 1 km upstream of bridge at Kordon Chernorechye, *Acer* & *Fagus* forest with *Sambucus*, *Corylus*, *Alnus*, *Ulmus* in understorey, 43.9319° N, 40.6778° E, 850 m a. s. l., 18 August 2012; 1 ♀ (CHR), Lagonakskiy Mountain Ridge, limestone cliffs on SW slope of Mt Matuk, 44.1061° N, 39.9225° E, 1810 m a. s. l., 16 August 2012; 2 ♀♀ (CHR), Mount Oshten, S of Lake Psheono-Dagh, limestone block field, 44.0092° N, 39.9008° E, 1970 m a. s. l., 25 August 2012, all leg. F. Walther; 1 ♂, 1 ♀ (SMNG), Mt Koryto, 44.067° N, 40.352° E, 1800 m a. s. l., August–September 1999, leg. O. Tietz; 4 ♂♂ (SMNG), Sakhray River, camp, 44°05'56" N, 40°23'56.4" E, 26 May 2004; 7 ♂♂, 6 ♀♀, 2 juv. (SMNG), W slope of Mt Asbestnaya, 44° 00' 28.8" N, 40° 27' 32.4" E, 26 August 2005; 1 ♀ (SMNG), W of Afonka Valley, 44° 00' 07.2" N, 40° 24' 21.6" E, 27 August 2005, all leg. K. Voigtlander; 5 ♂♂, 1 ♀, 3 juv. (ZMUM p2497). **Stavropol Prov.:** Zheleznovodsk, 10 July 1974; 4 juv. (ZMUM p2498), same locality, 29 June 1974; 2 ♂♂, 1 ♀, 6 juv. (ZMUM p2499), Zheleznovodsk, 15 July 1974; 2 ♂♂, 1 ♀, 1 juv. (ZMUM

p2500□□), same locality, foot of Mt Zheleznaya, deciduous forest, litter & under stones, 30 May 1982; 1 ♂ (ZMUM p2501), W of Zheleznovodsk, *Quercus* & *Crataegus* scrub, litter, 29 May 1982; 4 ♀♀, 1 ♂, 1 juv. (ZMUM p2502), 3 km E of Zheleznovodsk, forest of *Carpinus*, *Acer* etc., litter & under stones, 30 May 1982; 2 ♀♀ (ZMUM p2503), 3 km E of Zheleznovodsk, *Carpinus*, *Acer*, *Fraxinus* forest, litter & under stones, 30 May 1982; 4 ♂♂, 3 ♀♀ (ZMUM p2504), Pyatigorsk, Mt Mashuk, 600 m a. s. l., park of *Fraxinus*, *Acer*, *Quercus* etc., litter, 29 May 1982; 1 ♂, 3 ♀♀, 1 juv. (ZMUM p2505), E of Georgievsk, forest of *Quercus*, *Fagus*, *Acer*, litter & under bark, 7 June 1985; 1 ♂ (ZMUM p2506), same locality, 28 & 31 May 1982, all leg. S. Golovatch; 1 ♀ (ZMUM p2507), same locality, park in a sanatorium, litter, under stones, on walls, 1 June 1982; 1 ♂, 1 ♀ (ZMUM p2508), Stavropol, Russkiy Forest, 09 March 2014, leg. & det. R. Zuev; 1 ♂ (CHR), same locality, Russian Forest near Komsomolskiy Water Reservoir, deciduous forest (*Quercus*, *Carpinus*, *Acer*), camp, 45° 04' 81" N, 41° 95' 67" E, 530 m a. s. l., 22 August 2012, leg. F. Walther; 1 juv. (ZMUM p2509). **Kara-chaevo-Cherkessia:** Urup Distr., 3 km E of Pregradnaya, 800 m a. s. l., *Quercus*, *Fagus*, *Alnus*, *Corylus* etc. forest, litter & rotten wood, 3 August 1986; 2 ♂♂, 2 ♀♀, 7 juv. (ZMUM p2510), Lower Teberda S of Karachaevsk, 1000 m a. s. l., *Quercus*, *Fagus* etc. forest, litter, 3 August 1986; 1 ♂, 1 ♀ (ZMUM p2511), Teberda Nature Reserve, Mt Malaya Khatipara above Teberda Town, 29–30 May 1985; 1 ♀ (ZMUM p2512), Teberda Nature Reserve, Kizgich Canyon N of Arkhyz, wet riverine *Alnus* & *Betula* forest, 1450–1500 m a. s. l., litter, under bark & stones, 5 June 1985; 1 ♂ (ZMUM p2513), same locality, *Abies*, *Picea*, *Pinus*, *Fagus*, *Betula* & *Acer* forest, 1550–1650 m a. s. l., litter, under bark & stones, 5 June 1985; 1 ♀, 1 juv. (ZMUM p2514), Pass Gumbashi ca. 32 km NE of Karachaevsk, 2000 m a. s. l., subalpine meadow, under stones, 11 July 1986; 2 ♂♂ (ZMUM p2515), ca. 30 km S of Kurjinovo, Bolshaya Laba River valley, 4 km N of Damkhurts, 1050–1100 m a. s. l., *Fagus*, *Acer*, *Picea* etc. forest, litter & bark, 4 August 1986; 1 ♂, 1 ♀ (ZMUM p2516), Teberda Nature Reserve, Canyon Gonachkhir between Teberda & Dombai, road to Klukhor Pass, 1700–1900 m a. s. l., *Abies*, *Fagus*, *Acer* etc. forest, 1 June 1985; 2 juv. (ZMUM p2517), Teberda Nature Reserve, Mt Mussa-Achitara



Fig. 1. *Strongylosoma kordylamythrum* Attems, 1898, female from Adygea, habitus. A – dorsal view, B – lateral view. Photographed by A. Tikhonov, taken not to scale.




Fig. 2. *Strongylosoma kordylamythrum* Attems, 1898, male from Adygea. A – left gonopod, lateral view. B – left gonopod, lateral view. C – end part of left gonopod, ventral view. D and E – leg of the third pair. Photographed by V. Schmatko, scale bars: 2.0 mm. Del. A. Evsyukov, scale bars: 0.5 mm.

near Dombai, 2700–2800 m a. s. l., alpine meadow, under stones, 29 July 1986; 2 ♂♂, 5 ♀♀ (ZMUM p2518), Teberda Nature Reserve, Dombai, *Abies*, *Fagus*, *Picea*, *Betula*, *Acer* etc. forest, 1700–1800 m a. s. l., litter, under bark & stones, 31 May 1985, all leg. S. Golovatch; 1 ♀ (AE), Teberda Nature Reserve, Mt Malaya Khatipara, 24 July 2012, leg. D. Khisametdinova & Y. Kochetov; 1 ♂ (SMNG), “Kaukasus, Dombai, Cuchur-Tal, 2 June 1984, leg. W. Dunger”; 1 ♂ (ZMUC), 2–3 km NE of Dombai, SW slopes of Musat-Cheri Mt. Range, 43° 17'44" N, 41° 38' 24.7" E, 2000 m a. s. l., *Abies-Fagus-Picea* forest, sifting leaf litter 25 July 2011, leg. A. Solodovnikov; 2 ♂♂, 1 ♀, 1 juv. (ZMUM p2519). **Kabardino-Balkaria:** Baxan River valley, Bedyk between Tyrnyauz & Zhankhoteko, 900 m a. s. l., *Corylus*, *Fagus* & *Carpinus* scrub, litter & under stones, 20 July 1986; 1 ♂, 1 ♀, 5 juv. (ZMUM p2520), Lower Chegem, 800 m a. s. l.,

Fagus forest, litter, 14 July 1986; 1 ♂, 1 juv. (ZMUM p2521), Chegem Distr., 5 km S of Upper Chegem, 1700 m a. s. l., *Betula*, *Pinus* & *Juniperus* forest, litter, 12 July 1986; 6 ♂♂, 8 ♀♀, 2 juv. (ZMUM p2522), Chegem Distr., Upper Chegem, meadow, under stones, 1500 m a. s. l., 12 July 1986, all leg. S. Golovatch. 1 ♂, 6 ♀♀, 3 juv. (ZMUM p2523). **North Ossetia:** near Novo-Georgievskoye by Mozdok, *Quercus*, *Acer*, *Ulnus*, *Fraxinus* etc. forest, litter & under stones, under bark of logs, 27 May 1982; 1 ♀, 1 juv. (ZMUM p2524), S of Ordjonikidze (now Vladikavkaz), between Chmi & Baltik, *Quercus* & *Ulmus* forest on slope, litter & under stones, 2 June 1982, all leg. S. Golovatch; 1 ♂ (AE), Tanandon River valley, 2000 m a. s. l., forest, litter, 9 August 2010, leg. A. Evsyukov & D. Khisametdinova. **Abkhazia:** 1 ♂ (ZMUM p2525), environs of Sukhumi, Shromy, 30 April 1970; 1 ♂ (ZMUM p2526), same locality, 27 April 1970; 1 ♂, 1 ♀ (ZMUM p2527), "sample No. 8", under bark, 29 April 1971, all leg. N. Zalesskaya; 1 ♀ (ZMUM p2528), Sukhumi, 200 m a. s. l. SE of Besletsky Bridge, Cave Ammonalnaya I, 13 November 1987, leg. N. Myuge; 2 ♂♂, 2 ♀♀ (ZMUM p2529), Sukhumi, Botanical Garden, 9 October 1978; 1 ♂, 1 ♀ (ZMUM p2530), same locality, litter, 20 October 1978; 1 ♀ (ZMUM p2531), SE of Lake Ritsa, between Pass Anchkho & Pskhu, 1300–1450 m a. s. l., *Fagus*, *Acer* etc. forest on slope, under bark & stones, 14 & 16 August 1986; 1 ♂, 5 ♀♀ (ZMUM p2532), Myussera Nature Reserve, 20–130 m a. s. l., mixed deciduous forest (*Castanea*, *Alnus* etc.), litter, under bark & stones, 8–10 April 1983; 1 ♀ (ZMUM p2533), near Lake Ritsa, forest, litter, 24 October 1978; 2 ♂♂ (ZMUM p2534), Sukhumi Distr., Bzyb River valley, Pskhu, 700–950 m a. s. l., *Fagus*, *Quercus*, *Castanea* etc. forest, litter, under bark & stones, 15–16 August 1986; 6 ♂♂, 6 ♀♀ (ZMUM p2535), environs of Sukhumi, near Cave Kelassuri, litter, 11 April 1983; 1 ♂, 1 ♀ (ZMUM p2536), Lake Ritsa, 950–1100 m a. s. l., *Fagus*, *Abies*, *Picea*, *Acer* etc. forest, litter, under bark and stones, 13–14 August 1986, all leg. S. Golovatch; 6 ♂♂ (ZMUM p2537), Sukhumi Distr., Lower Yashtukha, 29 March 1985, leg. A. Markossian; 1 ♀ (ZMUM p2538), Tsandripsh, 15 August 2011, leg. E. Khatchikov. **Georgia:** 1 ♀ (ZMUM p2539), Ambrolauri Distr., near Nikortsminda, 1100 m a. s. l., near Cave Sakinule, 10 July 1974; 1 ♀ (ZMUM p2540), 40 km W of Mestia, Kherkhvashi E of Nakra, 1250–1700 m a. s. l., *Quercus*, *Fagus*, *Carpinus*, *Picea*, *Abies* etc. forest, litter & bark, 21 August–21 September 1986; 1 ♀ (ZMUM p2541), Borzhomi Distr., 8 km SE of Akhaldaba, 1000 m a. s. l., Nedzura River valley, *Picea*, *Carpinus* & *Fagus* forest, litter, logs, 12 May 1983, all leg. S. Golovatch; 2 ♀♀ (CHR), Samegrelo-Zemo Svaneti, Nodashi, 43° 03' 31.0" N, 42° 24' 59.0" E, 1100 m a. s. l., 29 September 2012, leg. F. Walther; 5 ♀♀ (CHR), Sairme Gorge, mixed forest, leaf litter (*Acer* & *Betula*), 41° 50' 48.5" N, 42° 48' 27.0" E, 2200 m a. s. l., 11 July 2013; 1 ♀ (CHR), Ambrolauri to Djava, E of Oni, NE of Pipileti, 42° 34' 04.4" N, 43° 29' 58.2 E, mixed forest, 1 July 2014, all leg. L. Mumladze. **Azerbaijan:** 1 ♂ (ZMUM p2542), ca 20 km E of Alazani River, September 1981, leg. S. Alekseev; 1 ♂ (ZMUM p2543), Mountainous Karabakh, ca. 15 km WSW of Mardakert, 1100 m a. s. l., *Quercus*, *Fagus*, *Acer* etc. forest, litter, 2 June 1987; 1 ♀ (ZMUM p2544), Mountainous Karabakh, Turshsu ca. 15 km S of Shusha, 1700 m a. s. l., *Quercus*, *Carpinus*, *Acer* etc. forest, litter. 3 June 1987, all leg. S. Golovatch & K. Eskov.

REMARKS. This species is very common and widespread throughout much of the Caucasus, also occurring in the adjacent Rostov-on-Don Prov. and Kalmykia (Evsyukov & Golovatch 2013). In the northern Caucasus (Stavropol Province), *S. kordylamythrum* has been found in bird nests (Golovatch & Matyukhin 2011), thus suggesting occasional ornithochory. The specimens in northernmost samples, both from the Rostov-on-Don Prov. and Kalmykia, tend to be considerably smaller than those from the Caucasus proper (Evsyukov & Golovatch 2013). Apparently, this reflects the less favourable conditions this species experiences at the northern periphery of its distribution. Obviously the same concerns the southernmost population that occurs in a city park in Tehran, Iran (Enghoff & Moravvej 2005).

Swarming was occasionally observed (Zheleznovodsk and Pyatigorsk, summer 1974, SG) (Golovatch & Matyukhin 2011).

Strongylotoma lenkoranum Attems, 1898 (Figs 3–5)

Strongylotoma lenkoranum Attems, 1898: 67: 314 (D).

Strongylotoma lenkoranum: Attems 1914: 227 (M), 1926: 247 (D), 1937: 38 (D); Verhoeff 1921: 42 (D), 1940: 37 (D); Lohmander 1936: 40 (D, R); Kobakhidze 1965: 391 (R); Jeekel 1968: 94 (M); Hoffman & Lohmander 1968: 84 (R); Samedov et al. 1972: 1245 (R); Rakhamanov 1972: 116 (R); Golovatch 1983: 169 (R), 1995: 127 (R); Enghoff & Moravvej 2005: 68 (R); Enghoff 2006: 190 (R); Nguyen & Sierwald 2013: 1268 (M).

Strongylotoma stragulatum Lohmander, 1932: 4 (D) – synonymized with *Strongylotoma lenkoranum* by Attems 1937: 38.

Strongylotoma leukoranum [lapsus calami!]: Lang 1964: 240 (R).

Strongylotoma lencoranum [lapsus calami!]: Bababekova 1996: 90 (R).



Fig. 3. *Strongylosoma lenkoranum* Attems, 1898 male from Azerbaijan, habitus. Same as in Fig. 1.

MATERIAL EXAMINED. **Armenia:** 4 ♂♂, 2 ♀♀ (ZMUM p2545), Kafan Distr., Shikahoh Nature Reserve, Shikahoh, 900–950 m a. s. l., *Quercus*, *Fagus*, *Carpinus* forest near spring, litter, logs, and under stones, 28 April 1983; 1 ♂, 1 ♀ (ZMUM p2546), Shikahoh Nature Reserve, Nerkin And, old *Platanus* stand along river, litter, in rotten wood, under stones, 600 m a. s. l., 30 April 1983; 2 ♂♂, 1 ♀ (ZMUM p2547), Megri Distr., above Kuris, 1500 m a. s. l., *Quercus* and *Acer* forest, litter, under bark & stones along springs, 26 April 1983; 3 ♂♂, 7 ♀♀ (ZMUM p2548), Yerevan, park, litter, 16 November 1985, all leg. S. Golovatch; 1 ♀ (ZMUM p2549), Stepanavan, 1600–1650 m a. s. l., *Quercus*, *Fagus*, *Carpinus* etc. forest, litter & under bark, 21–22 May 1987, leg. S. Golovatch & K. Eskov. **Azerbaijan:** 1 ♂, 1 ♀ (ZMUM p2550), ca. 5 km N of Kutkashen (now Gabala), 1150–1200 m a. s. l., *Fagus* & *Carpinus* forest, litter & rotten wood, 2 May 1987; 2 ♂♂ 1 ♀ (ZMUM p2551), Shemakha Distr., Pirkuli, near Observatory, 1200–1250 m a. s. l., *Quercus*, *Acer*, *Taxus* etc. forest, litter, 30 April 1987, all leg. S. Golovatch & K. Eskov; 1 ♂ (ZMUM p2552), same locality, 24 May 1988, leg. N. Loganova; 1 ♂, 5 ♀♀ (ZMUM p2553), SW of Kuba, 750 m a. s. l., *Fagus*, *Quercus*, *Carpinus* etc. forest, litter & under bark, 23 April 1987; 1 ♂, 2 ♀♀ (ZMUM p2554), ca. 12 km E of Ismailly, Girdyman-Chay River valley, 850–880 m a. s. l., *Fagus*, *Quercus*, *Carpinus*, *Acer* forest, litter & under bark, 1 May 1987; 2 ♂♂, 1 ♀, 5 juv. (ZMUM p2555), Mountainous Karabakh, Terter River valley, Nadirkhanly ca. 12 km NE of Kelbajar, 1200 m a. s. l., *Fraxinus* & *Juglans* stand,

litter, 1 June 1987, all leg. S. Golovatch & K. Eskov; 2 ♂♂, 1 ♀, 1 juv. (ZMUM p2556), Mountainous Karabakh, near Firyuza, Domy, 1650 m a. s. l., pine forest, 14 August 1977, leg. N. Zalesskaya; 4 ♂♂, 3 ♀♀ (ZMUM p2557), Masally Distr., Istsu ca. 8 km SW of Masally, *Quercus*, *Acer*, *Carpinus* etc. forest, 80–140 m a. s. l., litter, under bark & stones, 19–20 October 1983, leg. S. Golovatch.

REMARKS. This species is even more widely distributed than *S. kordylamythrum*, being known from Georgia, Azerbaijan, Armenia, Iran, and Turkey. It has also been recorded in Afghanistan (Golovatch 1995), where it is likely to have been introduced. In the Caucasus, it appears generally to be distributed more southerly and easterly than *S. kordylamythrum*.




Fig. 4. *Strongylosoma lenkoranum* Attems, 1898 male from Azerbaijan. Same as in Fig. 2.

CONCLUSION

Superficially, *S. kordylamythrum* and *S. lenkoranum* are very similar. Attems (1898) described them, based only on minor details of the conformation of the gonopods. It is not surprising, therefore, that they were sometimes confused (Verhoeff 1921) or misidentified (Muralewicz 1927, Lohmander 1932), until Lohmander (1936) found and illustrated sufficiently reliable characters for discriminating these two species. The diagnostic characters are lying in the structure of the gonopods, mainly the shapes of the distofemoral process and solenomere, and that of the ventral hump on male femur 3 (cf. Fig. 1 D and E and Fig. 2 D and E).

Our map (Fig. 5) shows that the distributions of *S. kordylamythrum* and *S. lenkoranum* in the Caucasus region differ considerably. Nguyen & Sierwald (2013) listed only “Georgia, Caucasus, Iran” for the former and “Georgia, Caucasus” for the latter. In fact, *S. kordylamythrum* occurs throughout the northern Caucasus (Krasnodar and Stavropol provinces, Adygea, Karachaev-Cherkessia, Kabardino-Balkaria, North Ossetia, Ingushetia, Chechenia and Dagestan), reaching



Fig. 5. Map showing the distributions of *Strongylosoma kordylamythrum* Attems, 1898 (asterisk) and *S. lenkoranum* Attems, 1898 (square).

both the Rostov-on-Don Region and Kalmykia in the north. In Transcaucasia, this species inhabits Abkhazia, Georgia and Azerbaijan, including Hyrcania both within southeastern Azerbaijan and northwestern Iran. In contrast, *S. lenkoranum* occurs only in Transcaucasia, including the eastern part of the Caucasus Major within Azerbaijan, most of the Caucasus Minor within Georgia (Kartli), Armenia and Azerbaijan (Mountainous Karabakh), and Hyrcania within both Azerbaijan and Iran. Moreover, this species is also known from northeastern Turkey and from Kabul, Afghanistan.

The distributions of these two species overlap only in two regions: Mountainous Karabakh and Hyrcania (Fig. 5). Both might seem to show parapatry, but in fact, based on the numerous new samples reported above, they are allopatric and never found together. The records of both species cited by Attems (1898) from near Lenkoran, Azerbaijan and by Lohmander (1932, 1936) on Mt Boghro-dagh, Iran, are not for syntopic, coexisting populations. Our Fig. 5 includes all known records of *S. kordylamythrum* and *S. lenkoranum* in the Caucasus and adjacent parts of Turkey and Iran.

As regards the patterns in the distributions of both *Strongylosoma* species in terms of altitude in the Caucasus, they are very similar in showing clear-cut preferences for either lowland or midmontane areas, mostly forest habitats below 2000 m a. s. l. Truly high-montane occurrences in (sub)alpine meadows are extremely rare. The same concerns troglophilic populations. It is this general similarity in habitat/ecological preferences that seems to account for their allopatry.

A c k n o w l e d g e m e n t s

We are most grateful to all colleagues who placed their material at our disposal. We are also greatly obliged to Vladimir Schmatko and Alexey Tikhonov, both of Rostov-on-Don, who skillfully took the pictures. Henrik Enghoff, ZMUC, kindly provided a reference to the only relevant sample in ZMUC.

R E F E R E N C E S

- ATTEMS C. 1898: System der Polydesmiden. I. Theil. *Denkschriften der Kaiserlichen Akademie der Wissenschaften zu Wien, Mathematisch-Naturwissenschaftliche Classe* **67**: 221–482.
- ATTEMS C. 1914: Die indo-australischen Myriopoden. *Archiv für Naturgeschichte* **80A** (4): 1–398.
- ATTEMS C. 1926: Etude sur les Myriopodes recueillis par M. Henri Gadeau de Kerville pendant son voyage zoologique en Syrie. *Voyage Zoologique d'Henri Gadeau de Kerville en Syrie* **1**: 221–266.
- ATTEMS C. 1937: Myriapoda 3. Polydesmoidea I. Fam. Strongylosomidae. *Das Tierreich* **68**: i–xxii, 1–300.
- BABABEKOVA L. A. 1996: Podtip trahejnye – Tracheata [The subphylum of tracheates – Tracheata]. Pp.: 89–97. In: ALIEV S. V. & KASYMOV E. G. (eds): *Životnyj mir Azerbajdzhana. Tom 2. Tip členistonogie [Animal Life of Azerbaijan. Volume 2. The Phylum of Arthropods]*. Baku: Elm, 412 pp (in Russian).
- ENGHOFF H. 2006: The millipedes of Turkey (Diplopoda). *Steenstrupia* **29**: 175–198.
- ENGHOFF H. & MORAVEJ S. A. 2005: A review of the millipede fauna of Iran. *Zoology in the Middle East* **35**: 61–72.
- EVSYUKOV A. P. & GOLOVATCH S. I. 2013: Millipedes (Diplopoda) from the Rostov-on-Don Region, southern Russia. *Arthropoda Selecta* **22**: 207–215.
- GHILAROV M. S. 1972: Počvenná fauna buroziomov Bol'sogo Kavkazskogo hrebla [Soil fauna of brown soils of the Greater Caucasus Range]. Pp.: 37–38. In: GHILAROV M. S. (ed.): *Problemy počvennoj zoologii. Materialy četvertogo vsesoúzного совещaniâ, Baku, 1972 [Problems in Soil Zoology. Materials of the 4th All-Union Conference, Baku, 1972]*. Moskva: Nauka, 176 pp (in Russian).
- GOLOVATCH S. I. 1983: A contribution to the millipede fauna of Iran (Diplopoda). *Annalen des Naturhistorischen Museums in Wien* **85B**: 157–169.
- GOLOVATCH S. I. 1995: On several new or poorly-known Oriental Paradoxosomatidae (Diplopoda Polydesmida), II. *Arthropoda Selecta* **3**: 127–137.
- GOLOVATCH S. I. & MATYUKHIN A. V. 2011: New records of millipedes (Diplopoda), mainly from bird nests, in European Russia. *Arthropoda Selecta* **20**: 115–116.
- HOFFMAN R. L. & LOHMANDER H. 1968: The Diplopoda of Turkey. *Mitteilungen des Hamburgischen Zoologischen Museums und Instituts* **65**: 61–121.
- JEEKEL C. A. W. 1968: On the classification and geographical distribution of the family Paradoxosomatidae (Diplopoda, Polydesmida). PhD Thesis. Rotterdam: Academisch Proefschrift, 168 pp (Privately Published).

- KOBAKHIDZE D. N. 1965: Spisok diplopod (Diplopoda) Gruzinskoy SSR [A list of the millipedes (Diplopoda) of the Georgian Soviet Republic]. *Fragmenta Faunistica* **11**: 390–398 (in Russian, with summaries in Polish and German).
- LANG J. 1959: K poznaniu mnogonožek (Diplopoda) territorii SSSR [To the knowledge of millipedes (Diplopoda) of the territory of the USSR]. *Zoologičeskij Žurnal* **38**: 1790–1796 (in Russian, with a summary in English).
- LANG J. 1964: Diplopoden aus Griechenland, Kreta, der Türkei und Iran aus dem Sammlungen von Dr. K. Lindberg. *Acta Societatis Zoologicae Bohemoslovacae* **28**: 237–246.
- LIGNAU N. G. 1903: Mnogonožki Černomorskogo poberež'ja Kavkaza. Die Myriopoden am Kaukasischen Schwarzmeerofer. *Mémoires de la Société des Naturalistes de la Nouvelle-Russie* **25**: 82–149 (in Russian and German).
- LIGNAU N. G. 1915: Vielfüßler aus Abchasiens. *Annuaire du Musée Zoologique de l'Académie Impériale des Sciences de Petrograd* **19**: 349–401.
- LJOVUSCHKIN S. I. & LOKSHINA I. E. 1975: K voprosu o svázi pešernych diplopod s počvou [To the question of the relationships of cave millipedes with the soil]. Pp.: 210–211. In: EITMINAVICIUTE I. (ed.): *Problemy počvennoj zoologii. Materialy pátogo vsesoúznnogo sovešaniâ, Vilnius, 1975* [Problems in Soil Zoology. Materials of the 5th All-Union Conference, Vilnius, 1975]. Vilnius: Institute of Zoology and Parasitology, Academy of Sciences of the Lithuanian Soviet Republic, 365 pp (in Russian).
- LOHMANDER H. 1932: Neue Diplopoden aus Persien. *Göteborgs Kungliga Vetenskaps- och Vitterhets-Samhälles Handlingar, Series 5B* **3**(2): 1–44.
- LOHMANDER H. 1936: Über die Diplopoden des Kaukasusgebietes. *Göteborgs Kungliga Vetenskaps- och Vitterhets-Samhälles Handlingar, Series 5B* **5**(1): 1–196.
- MURALEWICZ W. S. 1907: Zur Myriopodenfauna des Kaukasus. *Zoologischer Anzeiger* **31**: 329–351.
- MURALEWICZ W. S. 1911: Očerk fauny Myriopoda Kavkaza [Essay on the fauna of Myriopoda of the Caucasus]. *Mitteilungen des Kaukasischen Museums* **5**: 1–80 (in Russian).
- MURALEWICZ W. S. 1913: Členistonogiā. – Arthropoda. Glava VIII. Klass mnogonožki. – Myriapoda [Arthropods. – Arthropoda. Chapter VIII. Class of Myriapods. – Myriapoda]. *Trudy Obšestva Izuchenija Černomorskogo Poberež'ja, Sankt-Peterburg* **2**: 215–221 (in Russian).
- MURALEWICZ W. S. 1927: K faune Myriopoda Kavkaza [To the fauna of Myriopoda of the Caucasus]. *Acta Societatis Entomologicae Stauropolitanae* **3**(1): 1–7 (in Russian).
- RAKHMANOV R. R. 1972: Dvuparnonogie mnogonožki Lenkoranskoy zony Azerbajdžana [Millipedes of the Lenkoran zone of Azerbaijan]. P.: 116. In: GHILAROV M. S. (ed.): *Problemy počvennoj zoologii. Materialy četvertogo vsesoúznnogo sovešaniâ, Baku, 1972* [Problems in Soil Zoology. Materials of the 4th All-Union Conference, Baku, 1972]. Moskva: Nauka, 176 pp (in Russian).
- SAMEDOV N. G., BABABEKOVA L. A. & RAKHMANOV R. R. 1972: Razmešenie mnogonožek (Myriopoda) v različnyh tipah počv Lenkorani [Distribution of myriapods (Myriopoda) in various soil types of Lenkoran']. *Zoologičeskij Žurnal* **51**: 1244–1247 (in Russian, with a summary in English).
- TALIKADZE D. A. 1984: O faune dvuparnonogih mnogonožek (Diplopoda) Kolhidskoj provincii Kavkaza [On the millipede fauna (Diplopoda) of the Colchian Province of the Caucasus]. *Zoologičeskij Žurnal* **63**: 142–145 (in Russian, with a summary in English).
- NGUYEN A. D. & SIERWALD P. 2013: A worldwide catalog of the family Paradoxosomatidae Daday, 1889 (Diplopoda: Polydesmida). *Check List* **9**: 1132–1353.
- VERHOEFF K. W. 1921: Chilognathen-Studien (91. Diplopoden-Aufsatzz). *Archiv für Naturgeschichte* **86A**(12): 23–80.
- VERHOEFF K. W. 1940: Über Diplopoden aus der Türkei. Türkiyedeki diplopodlar hakkında. *Revue de la Faculté de Sciences de l'Université d'Istanbul, Séries B*, **5**(1/2): 1–49 (in German, with a summary in Turkish).
- ZUEV R. V. 2014: Preliminary data on the millipedes (Diplopoda) from the Stavropol Territory, northern Caucasus, Russia. *Arthropoda Selecta* **23**: 347–354.