

ACTA MYCOLOGICA

Vol. 47 (1): 97–107

2012

Myxomycetes of the Lipówka reserve in the Niepołomice Old Growth Forest (S Poland)

ANNA DROZDOWICZ¹, POLA SZOLC², ANNA BOCHYNEK¹, and AGNIESZKA SALAMAGA¹

¹ Institute of Botany, Jagiellonian University, Kopernika 27
PL-31-501 Kraków, anna.drozdowicz@uj.edu.pl

²14th Silesian Insurgents' Regiment School Complex, Szkolna 1
PL-44-300 Wodzisław Śląski, pszolc@poczta.onet.pl

Drozdowicz A., Szolc P., Bochynek A., Salamaga A.: *Myxomycetes of the Lipówka reserve in the Niepołomice Old Growth Forest (S Poland)*. Acta Mycol. 47 (1): 97–107, 2012.

Fifty eight species and three varieties of slime moulds were collected in the Lipówka reserve in 1999 and 2001. Thirty taxa of slime moulds very rare and rare in Poland were recorded in the reserve, including: *Arcyria minuta*, *A. stipata*, *Physarum penetrale*, *P. robustum*, and *Symphytocarpus flaccidus* which are red listed (Drozdowicz et al. 2006).

Key words: slime moulds, lignicolous species, biodiversity

INTRODUCTION

Myxomycetes are a small group of eukaryotic organisms with approximately 930 species described worldwide (Lado 2005-2012). About two hundred and fifty species have been recorded in Poland (Drozdowicz et al. 2003). Most data comes from national parks and nature reserves in Southern Poland, but in the recent years the interesting investigations were also undertaken in North-East part of the country (Panek, Romański 2010). The Lipówka reserve was selected as the study area on the quantitative and qualitative diversity of the myxomycete biota.

STUDY AREA

Situated in the northern part of the Niepolomice Old Growth Forest, the Lipówka reserve was established in 1957 and covers an area of 25.73 ha (Fig. 1). The reserve protects a natural, multi-species tree stand 170 to 200 years old. *Tilio-Carpinetum stachyetosum sylvaticae*, *Tilio-Carpinetum typicum*, *Circaeo-Alnetum* and *Carici elongatae-Alnetum* occur on moist and mesic habitats of the Vistula's alluvial floodplain. The flora of vascular plants comprises 134 species (Denisiuk 1978). This is consistent with the lowland forest flora of moist and shaded habitats. The tree stand in the reserve is formed by *Quercus robur*, *Carpinus betulus*, *Tilia cordata*, and *Alnus glutinosa*. *Pinus sylvestris* (planted), *Ulmus laevis*, and *U. minor* as well as sporadically *Padus avium* and *Populus tremula* occur as accompanying species.

A large amount of dead deciduous wood (logs, fallen trees, trunks and stumps, boughs and branches) was observed on the forest floor (Fig. 2). Bark was piled up in the area, especially around dead trunks of *Pinus sylvestris*. Thicker layers of litter were recorded in some parts of the reserve. They are potential microhabitats of slime moulds, but there were no information about occurrence of these organisms up to the 1999.

Macrofungi of Niepolomice Old Growth Forest were studied by Wojewoda (1978, 1979) and Komorowska (1980, 1986, 1991) but the most detailed observations were made in course of the project "Mycological monitoring in European oak forests" over the period 1994-1996 (Wojewoda et al. 1999).




Fig. 1. Geographical position of the Lipówka reserve (source: Nadleśnictwo Niepolomice).



Fig. 2. The „Lipówka” reserve – *Tilio-Carpinetum* in the early winter (05.01.2003; photo by R&M Kosiński).



Fig. 3. A common lignicolous slime moulds in the Lipówka reserve.
Lycogala epidendrum (L.) Fr. – immature aethalia and *Cribraria* sp. – black, globose, immature stalked sporangia (10.08.2010; photo by A. Bochynek).

MATERIAL AND METHODS

Research within slime moulds in the Lipówka reserve was carried out from July until November 1999 and from May until November 2001. The route method supplemented with observations of logs and stumps of selected tree species was used. Litter was also examined. Small samples of specimens were collected, i.e., single aethelia or aggregations of free sporangia, recorded at a locality. The substrate type, the collection date and frequency were noted for each taxon. A five-grade scale of the occurrence frequency by Drozdowicz (1992) was used.

The specimens were examined taxonomically using studies by Lister (1925), Krzemieniewska (1960), Martin and Alexopoulos (1969), Nannenga-Bremekamp (1991), Neubert, Nowotny and Baumann (1993, 1995, 2000), Stephenson and Stempfen (1994), Lado and Pando (1997), Ing (1999). The nomenclature was adopted after Lado (2005-2012), and vascular plants follows Mirek et al. (2002).

The collections of slime moulds were deposited in the Herbarium of the Institute of Botany, Jagiellonian University (KRA), in the MYXO division.

RESULTS

Sixty one taxa (58 species and 3 varieties) were identified among the total of 766 specimens collected. The nomenclatural information system of Eumycetozoa (Lado 2005-2012) does not include varieties while most taxonomical studies do. Specimens of typical forms of *Ceratiomyxa fruticulosa*, *Fuligo septica* and *Physarum viride* and their varieties were observed and collected in the Lipówka reserve. They are recorded on the list of slime moulds occurring in the reserve area.

Identified taxa are listed alphabetically and arranged in Table 1, the substrate categories colonised by the species, phenology and category of frequency are given for each taxon. Wood type is marked with an asterisk (*) if this is a new substrate supplementing the data for species reported by Drozdowicz et al. (2003). Based on research in Lipówka reserve, 39 species were noted on a substrate, hitherto not reported from Poland.

As it is indicated in Table 1, twenty two species occurred only on dead wood of lime, hornbeam and oak, two taxa were recorded on coniferous wood. Lignicolous species evidently dominated in the myxomycete biota and formed the most numerous ecological group of slime moulds in the Lipówka reserve. The greatest number of species, i.e. 32 were recorded on deciduous and coniferous trees, especially on logs lying on the forest floor and on logs suspended over drainage ditches. In this group, seven species of *Cribraria* were recognized in the Lipówka reserve. They occurred, first of all, on dead, deciduous wood. Only one of them, *Cribraria macrocarpa* was recorded on dead pine wood.

The following species were noted on the bark of living trees: *Lycogala epidendrum* (Fig. 3) on *Pinus sylvestris*, *Stemonitis fusca* on *Tilia cordata*, *Physarum album* and *P. cinereum* on *Carpinus betulus*. It could not be determined whether the presence of

Table 1
Biota of the slime moulds of the Lipówka reserve

No.	Species/varieties	Substrate type								Phenology		F	
		Qr	Cb	Tc	dw	P _s	cw	b	l	1999	2001		
1.	<i>Arcyria affinis</i> Rostaf.			+								08	I
2.	<i>Arcyria cinerea</i> (Bull.) Pers.	+	+	+	+	+	+	+				07-11	V
3.	<i>Arcyria denudata</i> (L.) Wettst.	+	+	+	+	+	+	+				05-11	IV
4.	<i>Arcyria ferruginea</i> Saut.	+	+	+	+	+	+	+				07-11	V
5.	<i>Arcyria minuta</i> Buchet		+	+		+						08	II
6.	<i>Arcyria obvelata</i> (Oeder) Onsberg	+	+	+		+						07-10	IV
7.	<i>Arcyria pomiformis</i> (Leers) Rostaf.	+	+	+		+						07-10	IV
8.	<i>Arcyria stipitata</i> (Schwein.) Lister	+	+	+		+						09-11	II
9.	<i>Ceratiumyx fruticulosa</i> (O. F. Müll.) T. Macbr. var. <i>fruticulosa</i>	+	+	+	+	+	+	+				05-10	V
10.	<i>Ceratiumyx fruticulosa</i> var. <i>porioides</i> (Alb. & Schwein.) G. Lister	+		+	+	+	+	+				07-08	II
11.	<i>Collaria arcyonema</i> (Racib.) G. Lister		+	+	+							08	I
12.	<i>Comatricha elegans</i> (Racib.) G. Lister		+	+	+							07-08	I
13.	<i>Comatricha nigra</i> (Pers. ex I. F. Gmel.) J. Schröt.	+	+	+		+						07-10	IV
14.	<i>Cribraria argillacea</i> (Pers. ex F. Gmel.) Pers.		+	+		+						07-10	II
15.	<i>Cribraria cancellata</i> (Batsch) Nann.-Bremek.				+		+					07	II
16.	<i>Cribraria macrocarpa</i> Schrad.					+						07	I
17.	<i>Cribraria piriformis</i> Schrad.	+	+	+		+	+					07-08	IV
18.	<i>Cribraria rufa</i> (Roth) Rostaf.	+	+	+		+	+					07-11	IV
19.	<i>Cribraria tenella</i> Schrad.				+		+					07-08	II
20.	<i>Cribraria vulgaris</i> Schrad.	+	+	+		+						07-08	III
21.	<i>Diachea leucopodia</i> (Bull.) Rostaf.			+								10	I
22.	<i>Dicydiaethalium plumbeum</i> (Schumach.) Rostaf.	+		+								07-08	IV
23.	<i>Diderma effusum</i> (Schwein.) Morgan		+	+								08	I
24.	<i>Diderma floriforme</i> (Bull.) Pers.		+	+								07-08	II
25.	<i>Diderma testaceum</i> (Schrad.) Pers.		+									07	I
26.	<i>Didymium clavus</i> (Alb. & Schwein.) Rabenh.		+									07	I
27.	<i>Didymium nigripes</i> (Link) Fr.											07	I
28.	<i>Fuligo leviderma</i> H. Neubert, Nowotny & K. Baumann	+	+	+	+	+	+	+				07-08	II
29.	<i>Fuligo septica</i> (L.) F. H. Wigg. var. <i>septica</i>	+	+	+		+						07-10	IV
30.	<i>Fuligo septica</i> var. <i>candida</i> (Pers.) R. E. Fr	+	+	+		+						07-10	IV
31.	<i>Hemirichia clavata</i> (Pers.) Rostaf.	+	+	+		+						05-11	V
32.	<i>Hemirichia sepulca</i> (Scop.) Rostaf. ex Lister	+	+	+		+						08-10	II

31.	<i>Lamproderma columbinum</i> (Pers.) Rostaf.																		09-11	I
32.	<i>Licea minima</i> Fr.																		07	II
33.	<i>Lycogala conicum</i> Pers.																		07-08	III
34.	<i>Lycogala epidendrum</i> (L.) Fr.																		05-11	V
35.	<i>Metatrichia floriformis</i> (Schwein.) Nann.-Bremek.																		10	II
36.	<i>Metatrichia vesparia</i> (Batsch) Nann.-Bremek.																		07-11	V
37.	<i>Physarum album</i> (Bull.) Chevall.																		07-11	V
38.	<i>Physarum cinereum</i> (Batsch) Pers.																		11	I
39.	<i>Physarum compressum</i> Alb. & Schwein.																		07; 10	II
40.	<i>Physarum contextum</i> (Pers.) Pers.																		08	I
41.	<i>Physarum globuliferum</i> (Bull.) Pers.																		07	II
42.	<i>Physarum leucophaeum</i> Fr.																		08-11	I
43.	<i>Physarum penetrale</i> Rex																		07-08	II
44.	<i>Physarum robustum</i> (Lister) Nann.-Bremek.																		07	I
45.	<i>Physarum viride</i> (Bull.) Pers. var. <i>viride</i>																		07-08	IV
	<i>Physarum viride</i> var. <i>aurantium</i> (Bull.) Lister																		07-11	III
46.	<i>Reitcularia lycoperdon</i> Bull.																		09-10	I
47.	<i>Stemonitis axifera</i> (Bull.) T. Macbr.																		05-11	IV
48.	<i>Stemonitis fusca</i> Roth																		07-10	IV
49.	<i>Stemonitopsis hyperopta</i> (Meyl.) Nann.-Bremek.																		07-11	III
50.	<i>Stemonitopsis typhina</i> (F. H. Wigg.) Nann.-Bremek.																		07-11	III
51.	<i>Symphycarpus flaccidus</i> (Lister) Ing & Nann.-Bremek.																		07-09	III
52.	<i>Trichia botrytis</i> (J. F. Gmel.) Pers.																		08	I
53.	<i>Trichia decipiens</i> (Pers.) T. Macbr.																		07-11	III
54.	<i>Trichia favoginea</i> (Batsch) Pers.																		07-11	V
55.	<i>Trichia persimilis</i> P. Karst.																		08-10	III
56.	<i>Trichia sabra</i> Rostaf.																		08-11	IV
57.	<i>Trichia varia</i> (Pers. ex J. F. Gmel.) Pers.																		09-11	IV
58.	<i>Tubifera ferruginosa</i> (Batsch) J. F. Gmel.																		05-08	III

Abbreviations: *Or* — *Quercus robur*; *Cb* — *Carpinus betulus*; *Tc* — *Tilia cordata*; *dw* — indeterminate deciduous wood; *Ps* — *Pinus sylvestris*; *cw* — indeterminate coniferous wood; *l* — litter; *F* — frequency; *I* — very rare species; *II* — rare species; *III* — frequent species; *IV* — very frequent species; *V* — common species; * wood type as a new substrate supplementing the data for species reported in “Myxomycetes of Poland. A checklist” (Drozdowicz et al. 2003).




Fig. 4. The occurrence of slime moulds in the Lipówka reserve in 1999 and 2001.

the sporangium of these slime moulds resulted from plasmodium migration from the trunk base or the development of the plasmodium *in situ* in bark fissures. Sporangia were situated on trunks at a height of 1 m to 2 m over the ground surface.

Diachea leucopodia, *Diderma testaceum*, *Didymium nigripes* and *Physarum contextum* were observed on litter, and one species *Diderma effusum* was recorded on oak wood and litter.

The intensity of the occurrence of individual myxomycete species varied in successive months of the vegetative seasons in 1999 and 2001 (Fig. 4). Specimens of species preserved from the previous vegetative season were mostly collected in May 2001, i.e., *Hemitrichia clavata*, *Metatrichia floriformis* and *Trichia favoginea*. The development of *Ceratiomyxa fruticulosa* and *Lycogala epidendrum* was also observed at that time.




Fig. 5. The frequency of slime moulds in the Lipówka reserve: I – very rare: 1–2 localities; II – rare: 3–5 loc.; III – frequent: 6–10 loc.; IV – very frequent: 11–29 loc.; V – common: > 30 loc.

The occurrence maxima of sporangia were observed in the Lipówka reserve in July and August in 1999 and 2001 when sporangia of up to 41 myxomycete species were recorded. Sporangia of 29 species were observed in September. Mature sporangia were also found in November. A considerable temperature drop inhibited plasmodium activity.

Two groups of taxa distinguished basing on the observations of slime moulds in two vegetative seasons with reference to the frequency scale (Fig. 5), were the most interesting: the groups I and II. Very rare myxomycetes (16 species) included species such as *Comatricha elegans*, *Physarum robustum* and *Symphytocarpus flaccidus*, rare slime moulds (13 species and 1 variety) included *Arcyria minuta*, *A. stipata* and *Physarum penetrale*. A total of 30 very rare and rare species were recorded in the reserve, and among them: *Arcyria minuta*, *A. stipata*, *Physarum penetrale*, *P. robustum* and *Symphytocarpus flaccidus* are rare in Poland according to the red list (Drozdowicz et al. 2006).

Lycogala conicum, *Stemonitopsis hyperopta* and other six species were frequent. Thirteen very frequent species were observed, including a.o.: *Arcyria pomiformis*, *Cribraria piriformis* and *Diderma effusum*. A total of eight species common in the reserve were found, including *Arcyria ferruginea*, *Ceratiomyxa fruticulosa* and *Lycogala epidendrum*.

DISCUSSION

Studies on slime moulds in the Lipówka reserve were conducted only in two vegetative seasons but their results are of great value as the myxomycete biota in the reserve had not been examined previously. Species rare in Poland were recorded, including *Arcyria minuta*, *A. stipata*, *Physarum penetrale*, *P. robustum* and *Symphytocarpus flaccidus* (Drozdowicz et al. 2006). The red list category of *Arcyria stipata* and *Symphytocarpus flaccidus* will probably change as the sites recorded in the Lipówka reserve are new localities of these species in Poland. New data on the colonisation of dead deciduous wood by individual myxomycete species collected in the present study expand the list of substrate types given in Myxomycetes of Poland (Drozdowicz et al. 2003).

The results of observations of myxomycetes in the Lipówka reserve confirm that there is no direct correlation between the occurrence of slime moulds and the plant community type. However, a multi-species tree stand with a high contribution of oak, hornbeam and lime indirectly influences the biota of myxomycetes. A variety of microhabitats of slime moulds have developed in the strictly protected part of reserve during the natural process of accumulation of high amounts of dead wood, mostly deciduous wood, on the forest floor. The degree of wood decomposition, the presence of bryophytes covering differently various parts of wood surfaces or bark fragments, often preserved on logs, branches and stumps, and the position of logs and branches on the soil surfaces affected the moisture content of substrata. Lignicolous taxa prevailed in the myxomycete biota. Only four species: *Diachea leucopodia*, *Diderma testaceum*, *Didymium nigripes*, *Physarum contextum* were recorded exclusively on the litter formed

by fallen leaves, fruits and seeds, cones, small twigs and bark fragments. Sporangia of *Lycogala epidendrum*, *Stemonitis fusca*, *Physarum album* and *P. cinereum* were observed on the bark of living trees; however, the species are not typical of corticolous myxomycetes (Ing 1994). In the event of *Cribraria* genus Krzemieniewska (1960) suggested that *Cribraria* spp., were most often found on coniferous wood, however the results of investigation in the Lipówka reserve were different – much more species of *Cribraria* were found on deciduous than on coniferous wood.

Slime moulds have been sometimes collected together with macromycetes during field studies in different forest complexes in Poland. In Central Poland slime

Table 2
Comparison of the slime moulds biota of the Lipówka and the Łęczok reserves

Taxa in both reserves	
<i>Arcyria cinerea</i>	<i>Lycogala conicum</i>
<i>Arcyria denudata</i>	<i>Lycogala epidendrum</i>
<i>Arcyria ferruginea</i>	<i>Metatrichia vesparia</i>
<i>Arcyria obvelata</i>	<i>Physarum album</i>
<i>Arcyria pomiformis</i>	<i>Physarum cinereum</i>
<i>Arcyria stipata</i>	<i>Physarum leucophaeum</i>
<i>Ceratiomyxa fruticulosa</i> var. <i>fruticulosa</i>	<i>Physarum viride</i> var. <i>viride</i>
<i>Comatricha nigra</i>	<i>Physarum viride</i> var. <i>aurantium</i>
<i>Cribraria piriformis</i>	<i>Reticularia lycoperdon</i>
<i>Cribraria rufa</i>	<i>Stemonitis axifera</i>
<i>Cribraria vulgaris</i>	<i>Stemonitis fusca</i>
<i>Diachea leucopodia</i>	<i>Stemonitopsis typhina</i>
<i>Dicthydiaethalium plumbeum</i>	<i>Trichia botrytis</i>
<i>Diderma effusum</i>	<i>Trichia favoginea</i>
<i>Diderma floriforme</i>	<i>Trichia persimilis</i>
<i>Fuligo leviderma</i>	<i>Trichia scabra</i>
<i>Fuligo septica</i> var. <i>septica</i>	<i>Trichia varia</i>
<i>Hemitrichia clavata</i>	<i>Tubifera ferruginosa</i>
<i>Hemitrichia serpula</i>	
Only in Lipówka reserve	
<i>Arcyria affinis</i>	<i>Fuligo septica</i> var. <i>candida</i>
<i>Arcyria minuta</i>	<i>Lamproderma columbinum</i>
<i>Ceratiomyxa fruticulosa</i> var. <i>porioides</i>	<i>Licea minima</i>
<i>Collaria arcyrionema</i>	<i>Metatrichia floriformis</i>
<i>Comatricha elegans</i>	<i>Physarum compressum</i>
<i>Cribraria argillacea</i>	<i>Physarum contextum</i>
<i>Cribraria cancellata</i>	<i>Physarum globuliferum</i>
<i>Cribraria macrocarpa</i>	<i>Physarum penetrabile</i>
<i>Cribraria tenella</i>	<i>Physarum robustum</i>
<i>Diderma testaceum</i>	<i>Stemonitopsis hyperopta</i>
<i>Didymium clavus</i>	<i>Symphytocarpus flaccidus</i>
<i>Didymium nigripes</i>	<i>Trichia decipiens</i> var. <i>decipiens</i>
Only in Łęczok reserve	
<i>Arcyria incarnata</i>	<i>Lamproderma arcyrionematis</i>
<i>Arcyria major</i>	<i>Lycogala exiguum</i>
<i>Arcyria oerstedii</i>	<i>Perichaena corticalis</i>
<i>Badhamia panicea</i>	<i>Perichaena depressa</i>
<i>Brefeldia maxima</i>	<i>Physarum flavicomum</i>
<i>Craterium aureum</i>	<i>Physarum pusillum</i>
<i>Craterium leucocephalum</i>	<i>Stemonitis flavogenita</i>
<i>Diderma deplanatum</i>	<i>Trichia affinis</i>
<i>Didymium iridis</i>	<i>Trichia contorta</i>
<i>Enerthenema papillatum</i>	<i>Trichia decipiens</i> var. <i>olivacea</i>
<i>Fuligo muscorum</i>	

moulds were collected in 7 nature reserves in *Tilio-Carpinetum* and *Potentillo albae-Quercetum* (Ławrynowicz 1973). This data are included in the list published by Kalinowska-Kucharska (1975). The most thorough investigations of slime moulds in *Tilio-Carpinetum* concentrated in the area of nature reserve Łęczczok (SW Poland) (Stojanowska 1974; Stojanowska, Panek 2002). The forest of this nature reserve comprises 136 ha and consists mainly of *Quercus robur*, *Carpinus betulus*, *Tilia cordata*, *Alnus glutinosa*, and *Ulmus glabra*, and some other trees, i.e., *Fagus sylvatica*, and *Picea abies*. Research on slime moulds in the Łęczczok reserve was conducted between 1967 and 1968, 1996 and 1999 and in 2001.

Thirty six species occur both in Łęczczok and Lipówka reserves. Twenty species and one variety were recorded in Łęczczok and twenty two species and two varieties were observed in Lipówka. The majority of taxa that occurred in both reserves are widespread and are common in Poland. However, *Arcyria stipata*, *Diderma effusum* and *D. floriforme* occur at few localities in Poland. Despite careful observations and a thorough in-field search, many taxa were recorded at single localities while many others were not found et all. The biotas recorded in both reserves are compared in Table 2.

It is difficult to identify specific factors that inhibit the occurrence of individual species. *Leocarpus fragilis* was not recorded either in the Lipówka reserve or in the Łęczczok reserve (Stojanowska, Panek 2002) while it is known to occur in southern Poland (Drozdowicz 1977, 1997). Although *L. fragilis* may be less numerous in the south than it is in the north of Poland (Panek, Romański 2010), it is recorded quite frequently mostly as a litter-colonising taxon.

CONCLUSIONS

Dead wood resources, specially deciduous wood and litter, constituted potential microhabitats for myxomycetes in both reserves. However, not only the quality and availability of the substrate is important for their development.

An overview of habitats and slime moulds associated with them worldwide was published in the late 20th century (Ing 1994). According to Ing, research into the occurrence, geographical distribution and relationships among various groups of organisms and slime moulds should be continued. Regional observations of slime moulds, conducted even in small areas, are important for a better understanding of these organisms.

REFERENCES

- Denisiuk Z. 1978. Szata roślinna rezerwatu Lipówka w Puszczy Niepołomickiej. Stud. Nat. A. 17: 87–117.
- Drozdowicz A. 1977. Myxomycetes of the Władysław Orkan forest reserve at Turbacz in the Gorce Mountains. Zesz. Nauk. Uniw. Jagiellon. Prace Bot. 5: 157–167 (in Polish with English summary).
- Drozdowicz A. 1992. Slime moulds (*Myxomycetes*) of the Ojców National Park. Part. I. Floristic problems. Zesz. Nauk. Uniw. Jagiellon. Prace Bot. 24: 125–145.
- Drozdowicz A. 1997. Studies on Myxomycetes in the Pieniny National Park I. New species for the PNP. Acta Mycol. 32 (2): 287–291.

- Drozdowicz A., Ronikier A., Stojanowska W., Panek E. 2003. Myxomycetes of Poland. A checklist. (In:) Z. Mirek (ed.). Biodiversity of Poland. 10. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Drozdowicz A., Ronikier A., Stojanowska W. 2006. Red list of rare Myxomycetes in Poland. (In:) Z. Mirek, K. Zarzycki, W. Wojewoda, Z. Szelać (eds). Red list of plants and fungi in Poland. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków: 91–99.
- Ing B. 1994. The phytosociology of myxomycetes. Tansley Review No 62. New Phytologist 126: 175–201.
- Ing B. 1999. The Myxomycetes of Britain and Ireland. The Richmond Publishing Co. Ltd. Slough, England.
- Kalinowska-Kucharska E. 1975. Materiały do flory śluzowców Polski Środkowej. Acta Mycol. 11 (1): 93–99.
- Kiszka J. 1978. Porosty rezerwatu Lipówka w Puszczy Niepołomickiej. Stud. Nat. A. 17: 149–158.
- Komorowska H. 1980. Mushrooms. (In:) K. Grodzińska (ed.). Acidification of forest environment (Niepołomice Forest) caused by SO² emissions from steel mills. Institute of Botany, Polish Academy of Sciences, Kraków: 91–99.
- Komorowska H. 1986. *Onygena equina* (Willd.) Pers.: Fr. in Poland. Acta Mycol. 22 (1): 49–52.
- Komorowska H. 1991. *Tricholomataceae* (Agaricales) Puszczy Niepołomickiej. Folia Soc. Sci. Lubl. 30, Biologia 1-2: 55–62.
- Krzemienievska H. 1960. Śluzowce Polski na tle flory śluzowców europejskich. Flora polska. PWN, Warszawa.
- Lado C. 2005–2012. An on line nomenclatural information system of Eumycetozoa. <http://www.nomen.eumycetozoa.com> (05.06.2012).
- Lado C., Pando F. 1997. Myxomycetes I. Ceratiomyxales, Echinosteliales, Liceales, Trichiales. Flora Mycol. Iberica 2: 1–323.
- Ławrynowicz M. 1973. Grzyby wyższe makroskopowe w grądach Polski środkowej. Acta Mycol. 9 (2): 133–204.
- Lister A. 1925. A monograph of the Mycetozoa. Ed. 3. London.
- Martin G. W., Alexopoulos C. J. 1969. The Myxomycetes. University of Iowa Press, Iowa City.
- Mirek Z., Piękoś-Mirek H., Zajac A., Zajac M. 2002. Flowering plants and pteridophytes of Poland. A checklist. (In:) Z. Mirek (ed.). Biodiversity of Poland. 1. W. Szafer Institute of Botany, Polish Academy of Sciences, Kraków.
- Nannenga-Bremekamp N. E. 1991. A guide to temperate myxomycetes. Biopress Limited, Bristol.
- Neubert H., Nowotny W., Baumann K. 1993. Die Myxomyceten Deutschlands und des angrenzenden Alpenraumes unter besonderer Berücksichtigung Österreichs. 1. Karlheinz Baumann Verlag, Gomaringen.
- Neubert H., Nowotny W., Baumann K. 1995. Die Myxomyceten Deutschlands und des angrenzenden Alpenraumes unter besonderer Berücksichtigung Österreichs. 2. Karlheinz Baumann Verlag, Gomaringen.
- Neubert H., Nowotny W., Baumann K. 2000. Die Myxomyceten Deutschlands und des angrenzenden Alpenraumes unter besonderer Berücksichtigung Österreichs. 3. Karlheinz Baumann Verlag, Gomaringen.
- Panek E., Romański M. 2010. Śluzowce *Myxomycetes*. (In:) L. Krzysztofiak (ed.). Śluzowce *Myxomycetes*, grzyby *Fungi* i mszaki *Bryophyta* Wigierskiego Parku Narodowego. Przyroda Wigierskiego Parku Narodowego, seria naukowa. Stowarzyszenie „Człowiek i Przyroda”. Suwałki: 9–85.
- Stephenson S. L., Stempen H. 1994. Myxomycetes. A handbook of slime molds. Timber Press, Portland, Oregon (Reprint 2000).
- Stojanowska W. 1974. Flora śluzowców kilku rezerwatów leśnych Opolszczyzny. Fragm. Flor. Geobot. 20 (2): 283–293.
- Stojanowska W., Panek E. 2002. Changes in the myxomycete biota of the „Łęczok” nature reserve near Racibórz (SW Poland). Acta Mycol. 37 (1/2): 13–28.
- Wojewoda W. 1978. Grzyby wielkoowocnikowe rezerwatu Lipówka w Puszczy Niepołomickiej. Stud. Nat. A. 17: 159–168.
- Wojewoda W. 1979. Rozmieszczenie geograficzne grzybów tremelloidalnych w Polsce. Acta Mycol. 15 (1): 75–144.
- Wojewoda W., Heinrich Z., Komorowska H. 1999. Macromycetes of oak hornbeam woods in the Niepołomice Forest near Kraków (S Poland) monitoring studies. Acta Mycol. 34 (2): 201–266.

Śluzowce rezerwatu Lipówka w Puszczy Niepołomickiej (Południowa Polska)

Streszczenie

Rezerwat Lipówka został utworzony w 1957 roku, dla zachowania starodrzewu *Tilio-Carpinetum stachyetosum sylvaticae*, *Tilio-Carpinetum typicum*, *Circaeo-Alnetum* oraz *Carici elongatae-Alnetum*. Uroczysko jest położone w północnej części Puszczy Niepołomickiej i zajmuje powierzchnię 25,73 ha (Ryc.1, 2). Na terenie rezerwatu Lipówka badano śluzowce w okresie od lipca do listopada 1999 roku oraz od maja do listopada 2001 roku. Stwierdzono występowanie 61 taksonów (58 gatunków i 3 odmiany) (Tab. 1). Większość śluzowców zanotowano na kłodach zarówno drzew liściastych, jak i iglastych (32 taksony). Na kłodach, pniakach, konarach i drobnych fragmentach martwego drewna drzew liściastych odnotowano 22 taksony, tylko 2 taksony występowały na drewnie drzew iglastych. Gatunki: *Diachea leucopodia*, *Diderma testaceum*, *Didymium nigripes*, *Physarum contextum* stwierdzono na ściółce. Natomiast zarodnie *Diderma effusum* zostały znalezione na martwym drewnie dębowym oraz na ściółce.

Śluzowce rozwijały się z różną intensywnością, a maksimum pojawów zaznaczyło się w lipcu i sierpniu (Ryc. 4). Odnotowano 29 taksonów bardzo rzadkich i rzadkich na obszarze uroczyska (Ryc. 5), wśród których do gatunków umieszczonych na Czerwonej liście śluzowców rzadkich w Polsce należą: *Arcyria minuta*, *A. stipata*, *Physarum penetrabile*, *P. robustum* oraz *Symphytocarpus flaccidus* (Drozdowicz et al. 2006).

W Polsce jednym z najlepiej opracowanych pod względem bioty *Myxomycetes* jest rezerwat Łęczczok (Stojanowska 1974; Stojanowska, Panek 2002), w którym lasy zajmują 136 ha. Jest to więc znacznie większy obszar niż uroczyska Lipówka, jednakże zaznacza się podobieństwo zbiorowisk leśnych wymienionych terenów. W rezerwacie Łęczczok badania nad śluzowcami przeprowadzono w latach 1967-68, 1996-99 oraz w 2001 roku.

Mimo opisanych różnic pod względem wielkości obszaru oraz długości okresów prowadzonych badań nad zróżnicowaniem gatunkowym śluzowców, dokonano porównania bioty śluzowców rezerwatów Łęczczok i Lipówka (Tab. 2). Lista gatunków wspólnych obejmuje 33 gatunki i 4 odmiany, natomiast tylko na obszarze rezerwatu Łęczczok znaleziono 20 gatunki i 1 odmianę, a w uroczysku Lipówka 21 gat. i 3 odmiany. Większość gatunków wspólnych, zebranych w obu rezerwach, należy do szeroko rozpowszechnionych i pospolitych na terenie Polski np. *Lycogala epidendrum* (Ryc. 3).