

Pupa with body tapering to a blunt point at the posterior extremity, which is provided with two short tubercles terminating in sharp spines. The head bears several stiff hairs; anterior and lateral margins of thorax bear row of stiff hairs; dorsal surface of thoracic and abdominal segments bears transverse row of stiff hairs; all hairs arising from tuberculate bases.

An Undescribed Species of Ptychoptera from the Western United States (Ptychopteridae, Diptera).

By CHARLES P. ALEXANDER, Urbana, Illinois.

The undescribed species of the genus *Ptychoptera* Meigen that is described herewith has been confused in collections with the much larger and darker *P. lenis* O. S.

Ptychoptera minor n. sp.

♂ Length, 9 mm.; wing, 8.6 mm. Rostrum and mouth-parts yellow; palpi yellow, the last segment dark brown. Antennae with the basal segment yellow, the second segment brown, the flagellum dark brown. Head black, shiny.

Pronotum yellow. Mesonotum black. Pleura black, gray pruinose, the dorso-pleural membranes bright yellow. Halteres yellow. Legs with the coxae yellow, marked on the outer face with purplish brown, the fore coxae with a basal band and an apical spot, the middle coxae with a basal and an apical band, the hind coxae with the outer face largely brown, only the apical margin yellowish; trochanters light yellow; femora yellow, the tips scarcely darkened; tibiae and metatarsi yellow, the tips darker; tarsi brown. Wings with a strong yellowish gray suffusion, the costal and subcostal cells and along vein *Cu* more yellowish; narrow, indistinct brown seams along the cord and at the fork of R_4+5 . Venation: *Sc* long, ending about opposite three-fourths the length of R_4+5 ; *Rs* moderately long, in alignment with R_4+5 , sometimes spurred at origin.

Abdominal tergites black, segments three to nine with the lateral margins broadly reddish, this color increasing in amount on the terminal segments, the hypopygium entirely reddish; sternites black, broadly ringed caudally with yellowish. Male hypopygium with the ninth tergite massive, deeply notched medially as in *P. lenis* but the lateral lobes only moderately slender, their apices widely separated, quite far before the tip on the ventral face with a small, blunt reddish lobe. Penis-guard not narrowed at the tip, the apex broadly rounded.

In *P. lenis* the hypopygium is black, the lobes of the ninth tergite slender, very narrowly separated at their apices, close before the tip



1920. "An undescribed species of Pilyhoptera from the western United States." *Entomological news, and proceedings of the Entomological Section of the Academy of Natural Sciences of Philadelphia* 31, 3-3.

View This Item Online: <https://www.biodiversitylibrary.org/item/84971>

Permalink: <https://www.biodiversitylibrary.org/partpdf/64110>

Holding Institution

University of Toronto - Gerstein Science Information Centre

Sponsored by

University of Toronto

Copyright & Reuse

Copyright Status: NOT_IN_COPYRIGHT

This document was created from content at the **Biodiversity Heritage Library**, the world's largest open access digital library for biodiversity literature and archives. Visit BHL at <https://www.biodiversitylibrary.org>.