

Polypores and associated beetles of the North Karelian Biosphere Reserve, eastern Finland

DMITRY S. SCHIGEL, TUOMO NIEMELÄ, MAARIT SIMILÄ, JUHA KINNUNEN and OLLI MANNINEN

SCHIGEL, D. S., NIEMELÄ, T., SIMILÄ, M., KINNUNEN, J. & MANNINEN, O. 2004: Polypores and associated beetles of the North Karelian Biosphere Reserve, eastern Finland. – Karstenia 44: 35–56. Helsinki. ISSN 0453-3402.

Polypores (poroid Basidiomycota) and associated beetles were inventoried in the Koitajoki Natura 2000 area (Ilomantsi municipality) and the Kitsi forest fire area (Lieksa), both belonging to the North Karelian Biosphere Reserve of the EU. As a result 105 polypore species were collected; together with earlier reports by other authors, the amount of species totals 121. This is a high number, surpassed in Finland by a few first-rank nature reserves only. Of the species found, 29 are red listed: 2 endangered (EN: *Antrodia crassa* (P. Karst.) Ryvarden, *Piloporia sajanensis* (Parmasto) Niemelä), 11 vulnerable (VU), and 16 near-threatened (NT). *Hyphodontia latitans* (Bourd. & Galz.) E. Langer has been found in Finland only once from the same area; now it was recollected. The research area is in a way a meeting-point of some northerly species (e.g. *Daedaleopsis septentrionalis* (P. Karst.) Niemelä, *Trichaptum laricinum* (P. Karst.) Ryvarden), those with an eastern distribution in Fennoscandia (e.g. *Trichaptum pargamenum* (Fr.) G. Cunn.), and some southern ones (e.g. *Pycnoporellus fulgens* (Fr.) Donk). Remarkably numerous were some species which indicate old growth forests of high conservation value (e.g. *Amylocystis lapponica* (Romell) Singer, *Antrodia albobrunnea* (Romell) Ryvarden, *A. crassa*, *Fomitopsis rosea* (Alb. & Schwein. : Fr.) P. Karst., *Phellinus nigrolimitatus* (Romell) Bourd. & Galz., *Skeletocutis stellae* (Pilát) Jean Keller). Beetle imagines were collected from polypore basidiocarps, and their larvae from basidiocarps and underlying decay, and then reared into adults. Special attention was paid to beetles living on rare polypore species. The polypore-associated beetle fauna totals 115 species, including 24 previously unrecorded from the Reserve. Our paper includes beetle records from ca. 30 such polypore species of which no previous beetle finds have been reported in the literature. The ecology of beetles living on fungal basidiocarps is discussed. Polypores can be divided into different ecological groups according to which beetles they attract; a division into *basidiocarp consistency classes* is proposed to describe such groups. Furthermore, the freshness or decomposition of a basidiocarp determines the amounts of beetles and their larvae, and their species composition.

Key words: *Basidiomycota*, *Coleoptera*, *Hyphodontia latitans*, *basidiocarp consistency class*, *beetles*, *Finland*, *polypores*, *ecology*

Dmitry S. Schigel, Tuomo Niemelä, Juha Kinnunen, Olli Manninen, Botanical Museum (Mycology), P. O. Box 7, FI-00014 University of Helsinki, Finland. Correspondence e-mail: dmitry.shchigel@helsinki.fi

Maarit Similä, Finnish Forest and Park Service, Natural Heritage Services, Urheilukatu 3 A, FI-81720 Lieksa, Finland.

1. Introduction

The Koitajoki river and its surroundings in the commune of Ilomantsi, Finnish North Karelia, make up the easternmost Natura 2000 site of the European Union. It is about 7400 ha wide and consists of protection areas with different protection statuses. There are the Koivusuo Strict Nature Reserve, protection areas for peatlands (Ruosmesuo – Hanhisuo and Ristisuo) as well as a reserve for old-growth forests. About 900 ha of the area are under restrictions of the Finnish forest law; there forest management operations are allowed outside of the important key habitats. The remaining 6500 ha belong to the Nature Reserve without forest management.

The Koitajoki Natura 2000 site is an important wilderness area in eastern Finland (Figs. 1, 2). It is situated in the transition zone of northern mires (pohjoiset aapasuot) and southern raised bogs (eteläiset keidassuot). In the area there are relatively large and natural mires and bogs with mosaic-like forests on mineral soil. Most of the forests are seminatural and old, with plenty of dead wood but with some old signs of previous selective cuttings. An important element of this Natura 2000 site is the Koitajoki, a shallow and meandering river with sandy banks (Figs. 3, 4). The Koitajoki Natura 2000 site maintains important rare and threatened species of polypores (Bondarceva et al. 1995; Bondartseva et al. 1998, 1999, 2001, Niemelä et al. 2002), beetles (Yakovlev et al. 2001) and other organisms.



Fig. 1. The research area in easternmost Finland, North Karelian Biosphere Reserve. The Jäkäläkangas Natura 2000 site is marked with an asterisk, rectangular box marks the Koitajoki Natura 2000 site, enlarged on Fig. 2.




Fig. 2. The Koitajoki Natura 2000 site, eastern Finland. Light grey covers the protected area, dark grey the Koivusuo Strict Nature Reserve. Black dots mark the centres of studied plots.

Fig. 3. Meanders of the Koitajoki in the North Karelian Biosphere Reserve near Polvikoski. Photograph TN.



Fig. 4. Old-growth forest in Liekkuaho, the Koitajoki Natura 2000 site. Photograph TN.



The Jäkäläkangas Natura 2000 site (224 ha) is situated in the municipality of Lieksa. Parts of the Jäkäläkangas area were burned by a wildfire in 1992. So the area includes about 143 ha of naturally burnt forests and peatlands which were protected immediately after the forest fire. About 17 ha were burnt completely to become a very interesting area with plenty of charred dead wood. Polypores were surveyed in the parts of less intensive fire.

Both areas surveyed in September – October 2002 and September 2003 are located inside the North Karelian Biosphere Reserve around Kitsi (63°16' N, 30°44' E). This first area of the Finnish Biosphere Reserves was established in 1992 (Hokkanen & Ieshko 1995). Biosphere Reserves are an essential part of UNESCO's Man and Biosphere (MAB) programme. They make up a backbone of the international network of research areas for sustainable development.

The inventory of the poroid Basidiomycete fungi (polypores) belongs to a multi-year study on Finnish polypores in various nature reserves, organized by the Finnish Forest and Park Service. It was started by Niemelä and Dai (1998, 1999) in the Luosto range, and continued in Korouoma (Niemelä & Kinnunen 2001), Ylläs – Aakenus (Niemelä & Kinnunen 2002) and Pisavaara (Niemelä & Kinnunen 2003), all of them in northern Finland. This paper is partly based on the 2002 inventory report (Niemelä et al. 2002), but additional collections were made in 2003, and our research materials were further worked out later. Data from the Repovesi National Park, Central Finland (Table 4) derive from an inventory in 2004 (unpublished). Although the polypores of Finland are rather well known, there is still a need in species inventories for better understanding of fungal ecology and nature protection.

European studies in the biology of saproxylic beetles started in the beginning of the 20th century in Scandinavia (Saalas 1917, 1923) and other parts of Europe (Donisthorpe 1931, 1935). From the 1950s the topic was favoured by many coleopterists (e.g. Benick 1952, Palm 1951, 1959, Pavour-Smith 1960). For an overview on Russian saproxylic beetles see Nikitsky and Schigel (2004).

Of more than 30 excellent papers published on the topic from 1990 onwards the most important ones by Økland (1995), Andersen et al. (2000), Thunes et al. (2000), Komonen et al. (2001), Martikainen (2001), Siitonen et al. (2001), Ehnström

and Axelsson (2002), Jonsell and Nordlander (2002), describe the fauna and ecology of Fennoscandian saproxylic beetles. Window and trunk beetle trappings (Yakovlev et al. 2001) showed the diversity of the saproxylic beetles of the Reserve. An emphasis of our study was put on search for previously unknown fungus – beetle interactions and details of their life cycles.

2. Materials and methods

Polypores and beetles were collected in forest compartments (geoinformation data of the Finnish Forest and Park Service). These compartments have been outlined so as to include fairly uniform stands of forest, and their areas and shapes vary according to the terrain.

In the field a complete list of polypore species was made from each forest compartment visited. These basic data were used to estimate the prevalences (frequencies of occurrence) of the species. The commonest, easily identified species were recorded *in situ*, but rare and difficult species and members of critical genera were collected for a closer scrutiny. Collected specimens were preliminarily studied in the microscope before drying. Collections were dried in mushroom dryers with ventilated air at +40–45°C. When needed, the identifications were later reconfirmed in laboratory with a research microscope, sections mounted in Cotton Blue or Melzer's reagent, and studied at $\times 1250$ magnification and phase contrast illumination. In addition to polypores, also other wood-inhabiting fungi were observed and collected, in particular rare and threatened species.


Phellinus igniarius is treated here in a wide sense (including *P. alni*, *P. cinereus*, *P. nigricans*), and *Postia lactea* and *P. tephroleuca* are listed collectively under the name *P. tephroleuca*.

Beetle imagines and larvae were collected separately and preserved in 70% alcohol for further identification. Ecological parameters were recorded in the field: basidiocarp hardness, which is both a characteristic of each polypore species and a result of decomposition process; moisture conditions; the presence of anamorphic fungi; decomposition stages of the basidiocarps (Thunes 1994). We used four visually defined successive stages of decomposition (Table 1, Figs. 5–8), omitting stage V in sampling, i. e. when detached basidiocarps are already dwelled by soil organisms. We would propose the use of subdivisions D (dry) and W (wet) for classes II–IV, while they were adopted by Thunes (1994) for the class IV only. Insect impact seems to be a result of insect attraction rather than pre-colonizing condition, so we modified the definitions of decomposition stages (Table 1).

Fruit bodies with beetle larvae were taken in plastic bags and boxes together with substrate for rearing, kept for 2–3 months in +4°C, and then exposed in room temperature for further two months before the checking of the rearing results. Breeding records meet the criteria set by Lawrence (1973: 165). The commonest Finnish polypores (Niemelä 2003a) with comparatively well known coleopteran fauna were not surveyed.

Table 1. Decomposition stages of polypore basidiocarps, according to Thunes (1994), modified. Subdivision D (dry) and W (wet) can be used for classes II–IV, if relevant.

Decomposition stage	Description
I	Alive, fresh and still actively growing basidiocarp.
II	Alive, fully grown, mature basidiocarp.
III	Dead, but fairly well-preserved basidiocarp; original outer and inner structures are still easily seen.
IV	Dead, strongly decomposed basidiocarp, structure transformed into unorganised mass.
V	Dead, detached and fallen basidiocarp, in transition of becoming incorporated in soil.



Figs. 5–8. Consecutive stages (I–IV) in the decomposition of the fruit body of *Haploporus odorus*. Photographs DSS, Pisavaara Strict Nature Reserve, northern Finland, 2003.

Fungal collections are preserved in the Herbarium of the Botanical Museum, Finnish Museum of Natural History, University of Helsinki (H); their nomenclature follows Niemelä (2004). Beetles will be donated after their investigation to the Zoological Museum, Finnish Museum of Natural History, University of Helsinki; single specimens have been given to identifiers (see Acknowledgements). Beetle nomenclature follows Silfverberg (1992), but the Ciidae are according to Müller et al. (2001). Authors of the Latin names of both fungi and beetles are given in Tables 2 and 3, respectively, and they are not repeated in the text.

All authors took part in the field work. This paper was prepared so that DSS wrote most of the text and all entomological sections. MS wrote parts of Introduction, and TN supervised the writing and wrote sections on fungal taxonomy.

3. Results

3.1 Polypores

Of the 105 polypore species collected by the authors, 29 are red-listed (Rassi et al. 2001) as belonging to different IUCN threat categories (2 EN – *Piloporia sajanensis*, *Antrodia crassa*; 11 VU; 16 NT; for details see Table 2). The list of species found by us is presented in Table 2, completed with some additional information. Our data to-

gether with finds published before (Bondarceva et al. 1995; Bondartseva et al. 1998, 1999, 2001) make a total of 121 polypore species known by now from the region of study.

3.2 Beetles

In our study 115 polypore-associated beetle species were found (Table 3), including 24 species previously unrecorded for the Reserve, in addition to the 91 known ones (Yakovlev et al. 2001). 2614 beetle adults and larvae were collected. No red-listed beetles were found or reared from polypores. Beetles or their larvae were found in 52 (49.5%) polypore species, while 53 polypore species appeared uninhabited (Table 2).

4. Discussion

4.1 Polypores and other fungi

The number of polypores found by us, 105 species (supplemented by 16 further species reported by others), is very high for any Finnish forest area. A recent inventory and herbarium study from the Pisavaara Strict Nature Reserve (Niemelä et

Table 2. Polypores (plus selected other Basidiomycota) and their beetles in the North Karelian Biosphere Reserve. Polypore species are given in alphabetical order. IUCN threat categories according to Rassi et al. (2001) specified after polypore names if present. N = number of compartments where found; A/B = number of fruit bodies inhabited/studied. Numbers of insect specimens found or reared are given in parentheses after the beetle species name. Imagines collected in nature indicated *light face*; insect records of larvae or reared imagines in *bold face*. Beetle – fungus associations meeting the Lawrence (1973) criterion are marked with “10+” instead of the number of larvae or reared imago specimens.

Fungus species	N	A/B	Insect records
1 <i>Amylocystis lapponica</i> (Romell) Singer VU	23	5/16	<i>Hapalarea linearis</i> (1), <i>Quedius xanthopus</i> (1 male), <i>Rhizophagus dispar</i> (1), <i>R. bipustulatus</i> (4), <i>Ostoma ferruginea</i> (2), <i>Dendrophagus crenatus</i> (1), <i>Cis comptus</i> (10+), <i>Hallomenus</i> sp. (10+ larvae)
2 <i>Anomoporia bombycina</i> (Fr.) Pouzar NT	3	0/3	
3 <i>Anomoporia kamtschatica</i> (Parmasto) M. Bondartseva	19	0/19	
4 <i>Antrodia albobrunnea</i> (Romell) Ryvarden NT	15	1/15	<i>Ostoma ferruginea</i> (3)
5 <i>Antrodia crassa</i> (P. Karst.) Ryvarden EN	1	0/1	
6 <i>Antrodia infirma</i> Renvall & Niemelä VU	4	0/4	
7 <i>Antrodia macra</i> (Sommerf.) Niemelä	6	0/6	
8 <i>Antrodia mellita</i> Niemelä & Penttilä VU	2	0/2	
9 <i>Antrodia primaeva</i> Renvall & Niemelä VU	2	0/2	

Fungus species	N	A/B	Insect records
10 <i>Antrodia pulvinascens</i> (Pilát) Niemelä VU	7	2/7	<i>Stenus carbonarius</i> (1 female), <i>Cis hispidus</i> (10+), <i>Dolichocis laricinus</i> (2) <i>Cis dentatus</i> (3)
11 <i>Antrodia serialis</i> (Fr.) Donk	43	1/8	
12 <i>Antrodia sinuosa</i> (Fr.) P. Karst.	42	0/42	
13 <i>Antrodia xantha</i> (Fr. : Fr.) Ryvarde	52	0/52	
14 <i>Antrodiella faginea</i> Vampola & Pouzar	2	0/2	
15 <i>Antrodiella semisupina</i> (Berk. & M.A. Curtis) Ryvarde	10	3/10	<i>Acrulia inflata</i> (3 males), <i>Rhizophagus dispar</i> (1), <i>Cis boletii</i> (1)
16 <i>Bjerkandera adusta</i> (Willd. : Fr.) P. Karst.	8	0/8	
17 <i>Byssoporia mollicula</i> (Bourdot) Larsen & Zak	6	0/6	
18 <i>Ceriporia viridans</i> (Berk. & Broome) Donk	2	0/2	
19 <i>Ceriporiopsis resinascens</i> (Romell) Domański	7	1/7	<i>Cis hispidus</i> (10+), <i>Octotemnus glabriculus</i> (2)
20 <i>Cerrena unicolor</i> (Bull. : Fr.) Murrill	17	1/17	<i>Dromius sigma</i> (1)
21 <i>Coltricia perennis</i> (L. : Fr.) Murrill	2	0/2	
22 <i>Daedaleopsis septentrionalis</i> (P. Karst.) Niemelä	1	0/1	
23 <i>Dichomitus squalens</i> (P. Karst.) D.A. Reid NT	8	2/8	<i>Scaphisoma boreale</i> (1), <i>Ennearthron cornutum</i> (10+)
24 <i>Diplomitoporus crustulinus</i> (Bres.) Domański NT	3	0/3	
25 <i>Diplomitoporus lindbladii</i> (Berk.) Gilb. & Ryvarde	4	0/4	
26 <i>Fomes fomentarius</i> (L. : Fr.) Fr.	59	1/1	<i>Bolitophagus reticulatus</i> (10+)
27 <i>Fomitopsis pinicola</i> (Sw. : Fr.) P. Karst.	57	3/57	<i>Ostoma ferruginea</i> (10+), <i>Rhizophagus dispar</i> (2), <i>Atomaria affinis</i> (2)
28 <i>Fomitopsis rosea</i> (Alb. & Schwein. : Fr.) P. Karst. NT	33	5/33	<i>Ostoma ferruginea</i> (4), Tineidae G. sp. (1)
29 <i>Ganoderma lipsiense</i> (Batsch) G.F. Atk.	6	0/6	
30 <i>Gelatoporia pannocincta</i> (Romell) Niemelä NT	9	4/9	<i>Rhizophagus dispar</i> (2), <i>Acrulia inflata</i> (1 male, 1 female), <i>Sepedophilus testaceus</i> (1 female), <i>Agathidium</i> sp. (1), Tineidae G. sp. (5)
31 <i>Gloeophyllum odoratum</i> (Wulfen : Fr.) Imazeki	1	0/1	
32 <i>Gloeophyllum sepiarium</i> (Wulfen : Fr.) P. Karst.	30	1/3	<i>Cis comptus</i> (9), <i>Sulcaxis affinis</i> (1)
33 <i>Gloeoporus dichrous</i> (Fr. : Fr.) Bres.	15	6/8	<i>Scaphisoma agaricinum</i> (1) <i>Cis comptus</i> (11)
34 <i>Gloeoporus taxicola</i> (Pers. : Fr.) Gilb. & Ryvarde	4	0/4	
35 <i>Hapalopilus rutilans</i> (Pers. : Fr.) P. Karst.	5	0/5	
36 <i>Heterobasidion parviporum</i> Niemelä & Korhonen	2	0/2	
37 <i>Hyphodontia latitans</i> (Bourd. & Galz.) E. Langer	1	0/1	
38 <i>Inonotus obliquus</i> (Pers. : Fr.) Pilát	50	0/50	<i>Rhizophagus bipustulatus</i> (1), <i>R. dispar</i> (1), <i>Dorcatoma dresdensis</i> (10+), <i>Triplax russica</i> (1), Elateridae G. sp. (1 dead larva) , <i>Abdera affinis</i> (2), <i>Mycetophagus quadripustulatus</i> (1)
39 <i>Inonotus radiatus</i> (Sowerby : Fr.) P. Karst.	2	0/2	<i>Abdera affinis</i> (3)
40 <i>Inonotus rheades</i> (Pers.) P. Karst.	11	5/11	<i>Corticaria rubripes</i> (1), <i>Epuraea variegata</i> (1), <i>Cis lineatocibratus</i> (10+), <i>Dorcatoma dresdensis</i> (1)
41 <i>Ischnoderma benzoinum</i> (Wahlenb.: Fr.) P. Karst.	11	0/11	
42 <i>Junghuhnia luteoalba</i> (P. Karst.) Ryvarde	13	1/13	<i>Sepedophilus testaceus</i> (1 male)
43 <i>Lenzites betulinus</i> (L. : Fr.) Fr.	1	1/1	<i>Dinaraea aequata</i> (1 male), <i>Cis hispidus</i> (10+)
44 <i>Leptoporus mollis</i> (Pers. : Fr.) Quél.	5	0/5	

Fungus species	N	A/B	Insect records
45 <i>Oligoporus rennyi</i> (Berk. & Broome) Donk	1	0/1	
46 <i>Oligoporus sericeomollis</i> (Romell) M. Bondartseva	25	0/25	
47 <i>Parmastomyces mollissimus</i> (Maire) Pouzar VU	1	0/1	
48 <i>Perenniporia subacida</i> (Peck) Donk NT	4	0/4	
49 <i>Phaeolus schweinitzii</i> (Fr.) Pat.	2	2/2	<i>Atheta boleticola</i> (1 male, 1 female), <i>Dorcatoma</i> sp. (6)
50 <i>Phellinus chrysoloma</i> (Fr.) Donk	29	5/7	<i>Abdera flexuosa</i> (1), <i>Ennearthron cornutum</i> (10+)
51 <i>Phellinus conchatus</i> (Pers. : Fr.) Quél.	13	4/13	<i>Rhizophagus dispar</i> (1), <i>Sulcaxis affinis</i> (2), <i>Cis hispidus</i> (2), <i>Dorcatoma</i> sp. (10+), Tineidae G. sp. (1)
52 <i>Phellinus ferrugineofuscus</i> (P. Karst.) Bourdot NT	27	1/4	Ciidae G. sp. (4 larvae)
53 <i>Phellinus igniarius</i> (L. : Fr.) Quél.	51	3/4	<i>Ennearthron cornutum</i> (10+) <i>Dorcatoma dresdensis</i> (10+)
54 <i>Phellinus laevigatus</i> (P. Karst.) Bourdot & Galzin	27	2/4	<i>Rhizophagus dispar</i> (4), <i>Acrulia inflata</i> (1 male), <i>Orthoperus atomus</i> (1), <i>Dorcatoma</i> sp. (3)
55 <i>Phellinus lundellii</i> Niemelä	21	2/21	<i>Ennearthron cornutum</i> (1), <i>Dorcatoma dresdensis</i> (3), Tineidae G. sp. (1)
56 <i>Phellinus nigrolimitatus</i> (Romell) Bourdot & Galzin	17	0/17	
57 <i>Phellinus pini</i> (Brot. : Fr.) A. Ames	40	6/14	<i>Phloeocharis subtilissima</i> (3 females + 2), <i>Hallomenus binotatus</i> (1), <i>Ennearthron cornutum</i> (10+) <i>Dorcatoma dresdensis</i> (10+)
58 <i>Phellinus populicola</i> Niemelä	10	0/10	<i>Dorcatoma dresdensis</i> (10+)
59 <i>Phellinus punctatus</i> (P. Karst.) Pilát	2	0/2	<i>Rhizophagus dispar</i> (1)
60 <i>Phellinus tremulae</i> (Bondartsev) Bondartsev & Borisov	44	0/5	
61 <i>Phellinus viticola</i> (Schwein. ex Fr.) Donk	45	1/7	<i>Cis boleti</i> (1), <i>Octotemnus glabriculus</i> (3), <i>Ennearthron cornutum</i> (10+)
62 <i>Physisporinus vitreus</i> (Pers. : Fr.) P. Karst.	2	0/2	
63 <i>Piloporia sajanensis</i> (Parmasto) Niemelä EN	1	0/1	
64 <i>Piptoporus betulinus</i> (Bull. : Fr.) P. Karst.	47	1/1	<i>Glischrochilus hortensis</i> (1), <i>Cis bidentatus</i> (10+), <i>Diaperis boleti</i> (10+)
65 <i>Polyporus brumalis</i> (Pers. : Fr.) Fr.	8	2/8	<i>Scaphisoma agaricinum</i> (1), Tineidae G. sp. (1)
66 <i>Polyporus ciliatus</i> Fr. : Fr.	3	0/3	<i>Rhizophagus dispar</i> (1)
67 <i>Polyporus leptcephalus</i> (Jacq. : Fr.) Fr.	10	3/10	<i>Orthoperus corticalis</i> (1), <i>Dolichocis laricinus</i> (1), <i>Cis jacquemartii</i> (2), <i>Cis lineatocribratus</i> (1)
68 <i>Postia alni</i> Niemelä & Vampola	15	2/15	<i>Hallomenus</i> sp. (2 larvae), Tineidae G. sp. (2)
69 <i>Postia caesia</i> (Schrad. : Fr.) P. Karst.	11	0/11	
70 <i>Postia fragilis</i> (Fr.) Jülich	5	2/5	<i>Hapalarea linearis</i> (1), <i>Hallomenus</i> sp. (2 larvae)
71 <i>Postia guttulata</i> (Peck) Jülich NT	7	0/7	
72 <i>Postia hibernica</i> (Berk. & Broome) Jülich NT	1	0/1	
73 <i>Postia lateritia</i> Renvall VU	6	3/6	<i>Hapalarea linearis</i> (1), <i>Lordithon lunulatus</i> (1 male), <i>Hallomenus ?binotatus</i> (2 larvae)
74 <i>Postia leucomallella</i> (Murrill) Jülich	19	3/19	<i>Hapalarea linearis</i> (1 male), <i>Rhizophagus dispar</i> (1), <i>Hallomenus</i> sp. (7 larvae)

Fungus species	N	A/B	Insect records
75 <i>Postia placenta</i> (Fr.) M.J. Larsen & Lombard NT	10	1/10	<i>Ischnoglossa prolixa</i> (1), <i>Rhizophagus dispar</i> (1), Hallomenus sp. (2 larvae)
76 <i>Postia septentrionalis</i> (Vampola) Renvall NT	1	0/1	
77 <i>Postia stiptica</i> (Pers. : Fr.) Jülich	2	1/2	Hallomenus sp. (2 larvae)
78 <i>Postia tephroleuca</i> (Fr.) Jülich	16	2/16	<i>Acrulia inflata</i> (1 female), <i>Atrecus pilicornis</i> (1), <i>Rhizophagus dispar</i> (3), Hallomenus binotatus (2), H. sp. (7 larvae)
79 <i>Postia undosa</i> (Peck) Jülich	4	0/4	
80 <i>Protomerulius caryae</i> (Schwein.) Ryvarde VU	5	0/5	
81 <i>Pycnoporellus fulgens</i> (Fr.) Donk	2	0/2	
82 <i>Pycnoporus cinnabarinus</i> (Jacq. : Fr.) P. Karst.	9	4/9	Sulcaxis affinis (11)
83 <i>Rigidoporus corticola</i> (Fr.) Pouzar	28	9/12	<i>Agathidium pisanum</i> (2), <i>Rhizophagus dispar</i> (1), <i>Acrulia inflata</i> (3 males, 1 female), Staphylinidae G. sp. (10+ larvae), Elateridae G. sp. (1)
84 <i>Sarcoporia salmonicolor</i> (Berk. & M.A. Curtis) Teixeira NT	2	1/2	Cis dentatus (6)
85 <i>Skeletocutis amorphia</i> (Fr.) Kotl. & Pouzar	13	0/13	
86 <i>Skeletocutis biguttulata</i> (Romell) Niemelä	14	0/14	
87 <i>Skeletocutis brevispora</i> Niemelä VU	6	0/6	
88 <i>Skeletocutis carneogrisea</i> A. David	4	0/4	
89 <i>Skeletocutis kuehneri</i> A. David	4	0/4	
90 <i>Skeletocutis lenis</i> (P. Karst.) Niemelä VU	2	0/2	
91 <i>Skeletocutis odora</i> (Sacc.) Ginns NT	6	1/6	Staphylinidae G. sp. (10+ larvae)
92 <i>Skeletocutis papyracea</i> A. David	5	0/5	
93 <i>Skeletocutis stellae</i> (Pilát) Jean Keller VU	4	0/4	
94 <i>Trametes hirsuta</i> (Wulfen : Fr.) Pilát	1	1/1	Cis hispidus (10+), Octotemnus glabriculus (10+)
95 <i>Trametes ochracea</i> (Pers.) Gilb. & Ryvarde	38	5/5	Cis boleti (10+), Cis hispidus (10+), Octotemnus glabriculus (10+), Tineidae G. sp. (1)
96 <i>Trametes pubescens</i> (Schumach. : Fr.) Pilát	10	4/4	Cis boleti (10+), Cis hispidus (10+)
97 <i>Trametes velutina</i> (Fr.) G. Cunn.	4	3/4	Cis boleti (10+), Octotemnus glabriculus (10+), Tineidae G. sp. (1)
98 <i>Trechispora candidissima</i> (Schwein.) Bondartsev	1	0/1	
99 <i>Trechispora hymenocystis</i> (Berk. & Broome) K.-H. Larsson	1	0/1	
100 <i>Trechispora mollusca</i> (Pers. : Fr.) Liberta	4	0/4	
101 <i>Trichaptum abietinum</i> (Pers. : Fr.) Ryvarde	53	0/53	
102 <i>Trichaptum fuscoviolaceum</i> (Ehrenb. : Fr.) Ryvarde	19	1/2	Cis punctulatus (10+)
103 <i>Trichaptum laricinum</i> (P. Karst.) Ryvarde NT	5	0/5	
104 <i>Trichaptum pargamenum</i> (Fr.) G. Cunn. NT	20	6/10	<i>Leptusa pulchella</i> (1), <i>Acrulia inflata</i> (1 male), <i>Rhizophagus dispar</i> (1), Cis comptus (8), Cis lineatocribratus (1), Ennearthron cornutum (2), Cis jacquemartii (2), Cis punctulatus (3)
105 <i>Tyromyces chioneus</i> (Fr.) P. Karst.	6	0/6	<i>Atheta</i> sp. (1)

Fungus species	N	A/B	Insect records
Non-polypore fungi			
106 <i>Amylostereum chailletii</i> (Pers. : Fr.) Boidin	1	0/1	
107 <i>Asterodon ferruginosus</i> Pat.	13	0/13	
108 <i>Basidioradulum radula</i> (Fr.) Nobles	5	0/5	
109 <i>Calocera cornea</i> (Batsch. : Fr.) Fr.	1		0/1
110 <i>Cantharellus tubaeformis</i> (Bull. : Fr.) Fr.	1	0/1	
111 <i>Chaetoderma luna</i> (Romell) Parmasto	8	0/1	
112 <i>Chondrostereum purpureum</i> (Pers. : Fr.) Pouzar	8	0/1	
113 <i>Columnocystis abietina</i> (Pers. : Fr.) Pouzar	14	0/14	
114 <i>Coniophora olivacea</i> (Pers. : Fr.) P. Karst.	13	0/13	
115 <i>Creolophus cirrhatus</i> (Pers. : Fr.) P. Karst.	2	0/2	
116 <i>Cydia salicina</i> (Fr.) Burt	1	0/1	
117 <i>Daldinia concentrica</i> (Bolton : Fr.) Ces. & De Not <i>s.l.</i>	1	0/1	
118 <i>Gloiodon strigosus</i> (Schwein. : Fr.) P. Karst. VU	3	0/3	
119 <i>Hericium coralloides</i> (Scop. : Fr.) Pers.	9	0/9	
120 <i>Hydnellum aurantiacum</i> (Batsch : Fr.) P. Karst.	1	0/1	
121 <i>Hydnellum ferrugineum</i> (Fr. : Fr.) P. Karst.	10	0/10	
122 <i>Hydnellum gracilipes</i> (P. Karst.) P. Karst.	1	0/1	
123 <i>Hypochnicium multiforme</i> (Berk. & Broome) Hjortst.	1	0/1	
124 <i>Hypsizygus ulmarius</i> (Bull.) Redhead	1	1/1	<i>Rhizophagus dispar</i> (10+)
125 <i>Kavinia albobiridis</i> (Morgan) Gilb. & Budington NT	1	0/1	
126 <i>Laeticorticium roseum</i> (Fr.) Donk	10	0/10	
127 <i>Laxitextum bicolor</i> (Pers. : Fr.) Lentz	3	0/3	
128 <i>Lentaria epichnoa</i> (Fr.) Corner	2	0/2	
129 <i>Lentinellus vulpinus</i> (Sowerby) Kühner & Maire	3	0/3	
130 <i>Mycena tintinabulum</i> Quéf. VU	1	0/1	
131 <i>Mycoacia fuscoatra</i> (Fr. : Fr.) Donk	7	0/7	
132 <i>Panellus serotinus</i> (Schrad. : Fr.) Kühner	1	0/1	
133 <i>Phanerochaete sanguinea</i> (Fr.) Pouzar	2	0/2	
134 <i>Phellodon niger</i> (Fr. : Fr.) P. Karst.	3	0/3	
135 <i>Phellodon tomentosus</i> (L. : Fr.) Banker	1	0/1	
136 <i>Phellodon secretus</i> Niemelä & Kinnunen	2	0/2	
137 <i>Phlebia centrifuga</i> P. Karst. VU	9	0/9	
138 <i>Phlebia cornea</i> (Bourd. & Galzin) Parmasto NT	2	0/2	
139 <i>Phlebia radiata</i> Fr.	3	0/3	
140 <i>Phlebia tremellosa</i> (Schrad. : Fr.) Burds. & Nakasone	18	0/18	
141 <i>Pholiota heteroclita</i> (Fr. : Fr.) Quéf.	1	0/1	
142 <i>Phyllotopsis nidulans</i> (Pers. : Fr.) Singer	3	0/3	
143 <i>Pleurotus dryinus</i> (Pers. : Fr.) P. Kumm.	1	1/1	<i>Rhizophagus dispar</i> (10+), <i>Sepedophilus testaceus</i> (3), <i>Hapalaraea melanocephala</i> (1) <i>Rhizophagus dispar</i> (10+), <i>R. bipustulatus</i> (10+), <i>Mycetophagus multipunctatus</i> (8), <i>Triplax aenea</i> (10+), <i>Atheta picipes</i> (10+), <i>Phloeopora testacea</i> (3), <i>Cerylon</i> sp. (7)
144 <i>Pleurotus pulmonarius</i> (Fr.) Quéf.	6	6/6	
145 <i>Plicatura nivea</i> (Sommerf. : Fr.) P. Karst.	9	0/9	
146 <i>Pseudohydnum gelatinosum</i> (Scop. : Fr.) P. Karst.	2		
147 <i>Pseudomerulius aureus</i> (Fr.) Jülich	6	0/6	
148 <i>Punctularia strigosozonata</i> (Schw.) Talbot	1	0/1	
149 <i>Sarcodon squamosus</i> (Schaeff.) Quéf.	3	0/3	
150 <i>Serpula himantoides</i> (Fr. : Fr.) P. Karst.	19	0/19	
151 <i>Sistotrema raduloides</i> (P. Karst.) Donk	10	0/10	
152 <i>Steccherinum ochraceum</i> (Pers.) Gray	1	0/1	
153 <i>Stereopsis vitellina</i> (Plowr.) D.A. Reid NT	1	0/1	

Fungus species	N	A/B	Insect records
154 <i>Stereum hirsutum</i> (Willd. : Fr.) Gray	11	0/11	
155 <i>Stereum rugosum</i> Pers. : Fr.	8	0/8	
156 <i>Stereum sanguinolentum</i> (Alb. & Schwein. : Fr.) Fr.	14	0/14	
157 <i>Thelephora terrestris</i> Ehrh. : Fr.	3	0/3	

Literature data, specimens not studied by us

Albatrellus confluens (Alb. & Schwein. : Fr.) Kotl. & Pouzar
Albatrellus ovinus (Schaeff. : Fr.) Kotl. & Pouzar
Albatrellus subrubescens (Murrill) Pouzar
Antrodiella romellii (Donk) Niemelä
Bjerkandera fumosa (Pers. : Fr.) P. Karst.
Ceriporia reticulata (H. Hoffm. : Fr.) Domański
Datronia mollis (Sommerf.) Donk
Glocophyllum protractum (Fr.) Imazeki
Haploporus odoratus (Sommerf.) Bondartsev & Singer
Hyphodontia flavipora (Cooke) Sheng H. Wu
Hyphodontia paradoxa (Schrad. : Fr.) E. Langer & Vesterholt
Junghuhnia collabens (Fr.) Ryvarden
Junghuhnia lacera (P. Karst.) Niemelä & Kinnunen
Polyporus badius (Pers.) Schwein.
Polyporus pseudobetulinus (Pilát) Thorn, Kotir. & Niemelä
Rigidoporus populinus (Schumach. : Fr.) Pouzar

al. 2004) yielded 125 species; that reserve is often considered to be the richest site in the whole of West and Middle Fennoscandian coniferous forest zone.

We have inventoried several forest reserves during recent years from different parts of North and East Finland (Niemelä & Dai 1998, Niemelä & Dai 1999, Niemelä & Kinnunen 2001, Niemelä & Kinnunen 2002, Niemelä et al. 2002, Niemelä & Kinnunen 2003). All they have been studied in the same way, by making full lists of polypore species from each forest compartment visited. The high number of compartments (593 compartments studied during the listed inventories, Table 4) enables us to make a summary on the commonest polypore species in these old and virgin forests. While most of the high-frequency species (*Fomes fomentarius*, *Trichaptum abietinum*, *Fomitopsis pinicola*, etc.) are able to inhabit many kinds of wooded biotopes, strikingly many are inhabitants of old-growth forests, and have virtually disappeared from areas where forest management has been practiced, for instance, tree stands thinned, dead trees removed, etc. This is a good example

on the impoverishing effect of modern forestry on forest biodiversity.

Our research site in the Koitajoki Reserve in eastern Finland lies at the transition between Middle Boreal and Southern Boreal zones (Ahti et al. 1968), in their slightly continental sections. The distributions of many northern species are known to reach further south in these climatically continental parts of East Fennoscandia, and similar results were obtained in our study, too. Among the polypores, *Daedaleopsis septentrionalis* and *Trichaptum laricinum* have typical northern distributions, and for instance in western Finland they are found almost exclusively within the Northern Boreal zone, i.e. in Lapland north of the Arctic Circle. However, they as well as some typically northerly hydneous fungi (*Hydnullum gracilipes*, *Phellodon secretus*; see notes below) and Corticiaceae (*Phlebia centrifuga*) were found in our research area, too.

Trichaptum parganenum has a clearly eastern distribution in Europe, and it is lacking from Central and West Finland, as well as from Sweden, Denmark and Norway (Hansen & Knudsen

Table 3. Systematic list of beetles attracted to polypores; North Karelian Biosphere Reserve. Numbers refer to host fungi (Table 2); light face = records of beetle imagines; **bold face** = larvae or rearings.

Taxon	New for the Reserve	Host fungi
Carabidae Latreille, 1802		
<i>Dromius sigma</i> (Rossi, 1790)	+	20(?)
Leiodidae Fleming, 1821		
<i>Agathidium arcticum</i> Thomson, 1862		<i>Reticularia</i> sp.
<i>Agathidium pisanum</i> Brisout de Barneville, 1872		83
Staphylinidae Latreille, 1802		
<i>Acrulia inflata</i> (Gyllenhal, 1813)	+	15, 30, 54, 78, 83, 104
<i>Atheta</i> (s.str.) <i>boleticola</i> J. Sahlberg, 1876	+	49
<i>Atheta</i> (<i>Traumoecia</i>) <i>picipes</i> (Thomson, 1856)	+	144
<i>Atheta</i> sp.	+	105
<i>Atrecus pilicornis</i> (Paykull, 1790)	+	78
<i>Dinaraea aequata</i> (Erichson, 1837)	+	43
<i>Hapalarea linearis</i> (Zetterstedt, 1828)	+	1, 70, 73, 74
<i>Hapalarea melanocephala</i> (Fabricius, 1787)	+	143
<i>Ischnoglossa prolixa</i> (Gravenhorst, 1802)	+	75
<i>Leptusa pulchella</i> (Mannerheim, 1830)	+	104
<i>Lordithon lunulatus</i> (Linnaeus, 1761)	+	73
<i>Quedius xanthopus</i> Erichson, 1839	+	1
<i>Phloeocharis subtilissima</i> Mannerheim, 1830	+	57
<i>Phloeopora testacea</i> (Mannerheim, 1830)	+	144
<i>Stenus carbonarius</i> Gyllenhal, 1827	+	10
<i>Scaphisoma agaricinum</i> (Linnaeus, 1758)		33, 65
<i>Scaphisoma boreale</i> Lundblad, 1952		23
<i>Sepedophilus testaceus</i> (Fabricius, 1792)	+	30, 42, 143
Elateridae Leach, 1815		
<i>G.</i> sp.		38, 83
Anobiidae Fleming, 1821		
<i>Dorcatoma dresdensis</i> Herbst, 1792		38, 40, 53, 55, 58
<i>Dorcatoma</i> sp.		49, 51, 54
Trogossitidae Latreille, 1802		
<i>Ostoma ferruginea</i> (Linnaeus, 1758)		1, 4, 27, 28
Nitidulidae Latreille, 1802		
<i>Eपुरaea variegata</i> (Herbst, 1793)		40
<i>Glischrochilus hortensis</i> (Goeffroy, 1785)		64
Monotomidae Laporte de Castelnau, 1840		
<i>Rhizophagus dispar</i> (Paykull, 1800)		1, 15, 27, 30, 38, 51, 54, 59, 66, 74, 75, 78, 83, 104, 124, 143, 144
<i>Rhizophagus bipustulatus</i> (Fabricius, 1792)		1, 38, 144
Cryptophagidae Latreille, 1802		
<i>Atomaria affinis</i> (F. Sahlberg, 1834)		27
Erotylidae Latreille, 1802		
<i>Triplax aenea</i> (Schaller, 1783)		144
<i>Triplax russica</i> (Linnaeus, 1758)		38
Cerylonidae Billberg, 1820		
<i>Cerylon</i> sp.		144
Corylophidae LeConte, 1852		
<i>Orthoperus atomus</i> (Gyllenhal, 1808)	+	54
<i>Orthoperus corticalis</i> (Redtenbacher, 1849)	+	67
Latridiidae Erichson, 1842		
<i>Corticaria rubripes</i> Mannerheim, 1844		40

Taxon	New for the Reserve	Host fungi
Ciidae Leach, 1819		
<i>Cis boleti</i> (Scopoli, 1763)		15, 61, 95–97
<i>Cis comptus</i> Gyllenhal, 1827		1, 32, 33, 104
<i>Cis dentatus</i> Mellié, 1848	+	11, 84
<i>Cis hispidus</i> (Paykull, 1798)		10, 19, 43, 51, 94–96
<i>Cis jacquemartii</i> Mellié, 1848		67, 104
<i>Cis lineatocribratus</i> Mellié, 1848		40, 67, 104
<i>Cis punctulatus</i> Gyllenhal, 1827		102, 104
<i>Cis bidentatus</i> (Olivier, 1790)		64
<i>Dolichocis laricinus</i> (Mellié, 1848)		10, 67
<i>Ennearthron cornutum</i> (Gyllenhal, 1827)		23, 50, 53, 55, 57, 61, 104
<i>Octotemnus glabriculus</i> (Gyllenhal, 1827)	+	19, 61, 94, 95, 97
<i>Sulcacis affinis</i> (Gyllenhal, 1827)	+	31, 51, 82
Mycetophagidae Leach, 1815		
<i>Mycetophagus quadripustulatus</i> (Linnaeus, 1761)		38
<i>Mycetophagus multipunctatus</i> Fabricius, 1792		144
Melandryidae Leach, 1815		
<i>Hallomenus</i> sp.		1, 68, 69, 74, 75, 77, 78
<i>Hallomenus binotatus</i> (Quensel, 1790)		57, 73, 78
<i>Abdera affinis</i> (Paykull, 1799)	+	38, 39
<i>Abdera flexuosa</i> (Paykull, 1799)		51
Tenebrionidae Latreille, 1802		
<i>Diaperis boleti</i> (Linnaeus, 1758)		64
<i>Bolitophagus reticulatus</i> (Linnaeus, 1767)		26

1997). In our research area it was found frequently on birch. *Piloporia sajanensis* seems to have a fairly continental, eastern distribution, too.

Some species of predominantly southern distribution were also found. *Pycnoporellus fulgens* is confined mostly to southern Fennoscandia (Niemelä 1980), but we recorded it once. This species has also been collected in Russian Karelia, not far from the Koitajoki area (Shubin & Krutov 1979), and hence its finding was not unexpected. Among the non-poroid Aphyllophorales with a southern distribution, we found *Punctularia strigosozonata*, *Steccherinum ochraceum*, *Stereopsis vitellina*, and among Agaricales *Mycena tintinabulum*, living on fallen trunk of birch.

Our records include a great number of threatened species: 2 endangered (EN), 11 vulnerable (VU), and 16 near-threatened (NT). They are indicated in the species list (Table 2). We made notes on the other wood-inhabiting fungi only in passing, when time allowed. Some threatened species

were found: *Mycena tintinabulum* (VU), *Gloiodon strigosus* (VU), *Kavinia alboviridis* (NT), *Phlebia centrifuga* (VU), *Phlebia cornea* (NT), *Punctularia strigosozonata* (CR), *Sistotrema raduloides* (NT) and *Stereopsis vitellina* (NT). These results clearly illustrate the high conservation values of the Koitajoki Natura 2000 site.

4.2 Notes on selected fungi

Antrodia crassa (Fig. 9) is extremely rare in Finland (Kotiranta & Niemelä 1996), and almost totally confined to the oldest pine forests. However, every now and then the species is found in more mesic, spruce dominated forests, indicating that dry environment *per se* is not obligatory for the species to grow. In our Koitajoki inventory the species was found in the oldest and best-preserved forest patch, the Kelokkoaho forest, which arises like an island in the middle of vast peatlands. Host tree was an exceptionally thick (over 50 cm), long-

ago fallen trunk of pine, on which also *Skeletocutis stellae* was growing. Long continuity of the forest, undisturbed conditions, and a very old, thick, fallen kelo tree (see Niemelä et al. 2002) seem to be needed for *A. crassa* to survive.

Piloporia sajanensis (Fig. 10) is a rarity throughout its range. It belongs to the so-called successor species (Niemelä et al. 1995; see below), which mostly inhabit trees that have first been decayed by other fungi. Such successors are fairly specific in terms of their preceding species; *Piloporia sajanensis* lives almost exclusively on trees decayed by *Trichaptum larici-*

num or, more seldom, the other *Trichaptum* species. Our sole find of *P. sajanensis* was growing on a fallen trunk of spruce, effectively white-rotted by *T. laricinum*, whose basidiocarps emerged in hundreds along the whole trunk. There are about 10 records of this vulnerable species in Finland, all of them from northern or easternmost parts of the country.

Hyphodontia latitans was reported from the same area already by Bondartseva et al. (1998). We recollected the species in another site of the Koitajoki Reserve, from a thin (9 cm diam.), strongly decayed fragment of spruce trunk. These are



Fig. 9. Fruit body of *Antrodia crassa*. Posio, Korouoma Nature Reserve, 2001, Niemelä 7085.



Fig. 10. Fruit body of *Piloporia sajanensis*. North Karelian Biosphere Reserve, Tapionaho, 2002, Niemelä 7496.

the only records of the species in Finland. The material was kindly identified by Heikki Kotiranta (Helsinki) and Karl-Henrik Larsson (Göteborg).

Hydnellum gracilipes is a rare, northern species of hydnaceous fungi. It was thoroughly described, illustrated and discussed by Kõljalg and Renvall (2000); at that time it was known from seven sites in Finland and one from Norway. Recent inventories have revealed a handful of new localities, but anyhow the species is very rare. All the collections were made in old, dry pine woodlands, where the basidiocarps of *H. gracilipes* are found growing in the small space between long-ago fallen kelo trunk and forest soil. The fragile, rhizomorph-like stipe arises from the ground, and the pileus spreads along the wood surface above.

Phellodon secretus was described recently (Niemelä et al. 2003), almost exclusively from the materials of our inventories. This slender, pale ash-grey species resembles *Phellodon connatus* (Schultz : Fr.) P. Karst., but its spores are smaller and context hyphae make a soft and loose, interwoven structure, while *P. connatus* has densely packed and parallel contextual hyphae. The holotype of this species is one of our Koitajoki collections. Both this and *H. gracilipes* are surely threatened, but their Red List statuses in Finland have not yet been established.

Punctularia strigosozonata is a stereoid fungus, characterized by soft, small, cigar brown pilei and usually an effused-reflexed habit of the basidiocarps. Bondartseva et al. (2000) made a detailed overview on the species in Europe, where its distribution is clearly eastern, continental. Now this rare species has been found also in Finland: one record from our inventory area, plus two others nearby, close to the Mekrijärvi Biological Station of the University of Joensuu in Ilomantsi, and an old collection by TN from the Koli National Park in Lieksa. All these were found growing on *Populus tremula*. First Finnish finds were reported by Niemelä (2003b).

4.3 Basidiocarp consistency classes

It is no surprise that in the North Karelian Biosphere Reserve both the species composition of beetles and their spatial distribution inside polypore fruit bodies appeared characteristic for taiga zone in general. Up to 73% of fungivore beetles are known to be polyphagous (Schigel 2002).

They often show no preference to certain polypore genera or groups of related genera, but colonize polypores of certain *consistency classes* (Table 5).

This concept of *basidiocarp consistency classes* is here proposed to be used when describing different kinds of fungi as habitats for insects and their larvae. It is not yet fully understood, which particular characteristics of fungal basidiocarps are ecologically decisive to make them suitable for beetles. Critical are, for instance:

- Shape and volume of the fruit body;
- Annuality vs. perenniality, and how long the perennial ones persist;
- Presence or absence of certain structures, e.g. crust;
- Water contents of the basidiocarp;
- Toughness of the mycelium, which depends on the hyphal system (monomitic, dimitic, trimitic), thickness of hyphal walls, and how dense the structure is;
- Chemical characteristics of the fruit body.

The division of basidiocarp consistency classes, proposed here, is based on an informal classification long used by mycologists while describing species and genera. Consequently, the names of the consistency classes derive from certain polypore genera, but here taxonomy is omitted and the terms represent patterns of physical characteristics only. Basidiocarp consistency classes are outlined in Table 5.

Although these consistency classes seem to be valid and are repeatedly found in nature, sometimes it is difficult to define sharp borders between them: for instance, *Ischnoderma resinosum* Fr. (P. Karst.), a central European species, is tyromycetoid (= leptoporoid) when young but turns fomitoid when old (Pouzar 1971). Another example of uncertainty is found in the genus *Trichaptum*: *T. abietinum*, *T. fuscoviolaceum* and possibly *T. laricinum* (more data needed), all growing on coniferous trees, have a characteristic set of beetle species, unlike *T. pargamenum* on birch and other deciduous trees, which is usually colonized by specialists of the trametoid consistency class. Beetles of resupinate polypores are much less well known, and these fungi are more difficult to be addressed in certain consistency classes. Anyhow, some of them could well be placed in the same classes as the pileate ones, for instance *Postia placenta* and *Sarcoporia salmonicolor* among the tyromycetoid ones.

Table 5. Basidiocarp consistency classes, and examples of characteristic polypore and beetle genera in European southern taiga.

Fomitoid

Hard, perennial, voluminous fruit bodies with thick context and several annual layers in hymenophore; robust when living; basidiocarps may stay attached on substrate for several years after death.

Examples of fungal genera: *Fomes*, *Fomitopsis*, *Phellinus*, *Ganoderma*, *Heterobasidion*.

Examples of beetles associated: *Bolitophagus*, *Ennearthron*, *Dorcatoma*, *Oplocephala*, *Ropalodontus*, *Cis* (subg. *Eridaulus*).

NB: In addition to beetle larvae adapted to develop in the hard context and trama, these fungi sometimes attract untypical imago, visiting sporulating fruit bodies (Latridiidae), or those covered by slime moulds (Leiodidae) or anamorphic fungi (Cryptophagidae). Larvae of beetles occur mostly in dying or dead fruit bodies. The life cycle may take more than one year.

Trametoid

Corky or leathery, projecting and fairly thin, shelf-shaped, fairly quickly drying, trimitic fruit bodies, annual.

Examples of fungal genera: *Trametes*, *Daedaleopsis*, *Funalia*, *Lenzites*, *Gloeoporus*, *Pycnoporus*, *Cerrena*, *Bjerkandera*.

Examples of beetles associated: *Cis* (*C. comptus*, *C. hispidus*, *C. micans*), *Sulcacis*, *Octotemnus*, *Tritoma*, *Wagaicis*.

NB: The tough context of dead fruit bodies is usually effectively eaten (generally by Ciidae) in dry condition the next season after sporulation. Larvae start to develop in living or dying fruit bodies. Several generations may utilize the cluster of fruit bodies before it is completely eaten.

Tyromycetoid

Soft and watery, monomitic, annual fruit bodies.

Examples of fungal genera: *Tyromyces*, *Postia*, *Amylocystis*, *Leptoporus*, *Hapalopilus*, *Pycnoporellus*.

Examples of beetles associated: *Hallomenus*.

NB: The high moisture contents of the fruit body and its short persistence limit the number of beetle species. Only two species of *Hallomenus*, pupating in soil, were found. Larvae eat living fruit bodies. Larval development is fast, one generation per year.

Piptoporoid

Corky or fleshy, voluminous, di/trimitic fruit bodies with thick and homogeneous context.

Examples of fungal genera: *Piptoporus*, *Polyporus*, *Laetiporus*.

Examples of beetles associated: *Mycetophagus*, *Diaperis*, *Dacne*, *Eledona*.

NB: Usually relatively large and thick fruit bodies, hosting both surface- and context-living larvae, which start to develop in either living or dying fruit bodies.

Xanthochroic

Brown coloured and monomitic, annual, at first fibrous, but becoming brittle upon dying.

Examples of fungal genera: *Inonotus*, *Onnia*.

Examples of beetles associated: *Abdera*, *Orchesia*, *Mycetophagus*.

NB: Larvae occupy the context of mostly living fruit bodies in somewhat similar way as the trametoid ones. Larvae very seldom pupate inside the fruit bodies, even if those usually stay on trunks for one or more years after their death. Beetles almost never colonise dead fruit bodies.

Trichaptoid

Thin, numerous, dimitic fruit bodies with purple coloured hymenophore; pilei merging at bases, annual, or continuing to grow over the next year.

Examples of fungal genera: *Trichaptum* (*T. abietinum*, *T. fuscoviolaceum*, ?*T. laricinum*).

Examples of beetles associated: *Cis* (*C. punctulatus*), *Wanachia*, *Zilora*.

NB: Beetle larvae settle at the confluent bases of the fruit bodies, where context thickness is sufficient to host larvae (Schigel 2002).

Several polypore genera like *Climacocystis*, *Gloeophyllum* and *Fistulina* (Nikitsky & Schigel 2004) have characteristic species assemblages and make consistency classes of their own. We see no sense to construct special names for them as far as there are no other genera sharing the same ecological characteristics.

4.4 Families of fungivorous beetles

Our study revealed a typical (Schigel 2002, Nikitsky & Schigel 2004) palearctic set of beetle families that are linked to polypores, among which the Ciidae, Anobiidae (*Dorcatoma*), Melandryidae and Tenebrionidae are the most efficient basidiocarp destructors and decomposers. The Staphylinidae, Nitidulidae, Leiodidae, Trogossitidae, Latridiidae (*Corticaria*) and Corylophidae (*Orthoperus*) visit polypores as adult beetles. They feed on various parts of fruit bodies or secondary organisms, such as anamorphic fungi or slime moulds covering dead basidiocarps. The highest number of species were found among imaginal visitors, although the less diverse Ciidae, Anobiidae, Melandryidae and Tenebrionidae tend to be more abundant and were found either as larvae or were reared (Tables 2–3).

Fungivorous beetles of different families utilize fungal basidiocarps in different ways. Slime mould specialists in Leiodidae often visit polypores during the sporulation period together with Corylophidae and Latridiidae. The most diverse family, Staphylinidae, contains just a few proved fungivorous species, although many species recorded as imagines usually visit moist (both living and dead) polypores as well as agarics and boletes. Strongly decomposed and wet fruit bodies attract saprophagous imagines of the Silphidae, Hydrophilidae, Cholevidae, and Scarabaeidae. Anobiidae (*Dorcatominae*) larvae develop in the hardest polypores of the fomitoid consistency class. Trogossitidae larvae feed on wood-rotting mycelium, but imagines often stay on polypore hymenophore. Cryptophagidae use such anamorphic fungi that cover dead polypores. Erotylidae live on wood-rotting fungi and have rather short life cycles, which allows them to use ephemeral *Pleurotus* species and some short-persisting polypores (*Inonotus obliquus*, *Trametes* spp., *Polyporus* spp., *Piptoporus betulinus*). Ciidae live in various polypores which stay dry at least part of the decomposition time. Mycetophagidae,

Tetratomidae and Melandryidae larvae with short cycles occupy annual tyromycetoid and xanthochroic polypores. Each Tenebrionidae species colonizes a narrow set of polypore species, while the family as a whole has a fairly wide ecological amplitude, decomposing polypores of various consistency classes.

4.5 The role of beetles as vectors of successor polypore species

Niemelä et al. (1995) dealt with a very special link in which certain rare polypores share the woody substrate with a number of common poroid or hymenochaetoid fungi. In these cases a common and effective decayer inhabits a tree trunk. When its mycelium becomes senescent, another fungus species invades the tree, possibly killing the mycelium of the first species, and often fruiting on its dead basidiocarps. These predecessor–successor links may be fairly common in boreal forests, but little is known on their ecological background and even less on the mechanisms how the trees inhabited by a predecessor become inoculated by the mycelium of a successor.

By the time when these predecessor–successor relations were described, not much was understood about the spore dispersal of successor species, which are usually fairly rare and selective about their predecessors. We found that certain beetle larvae and successor polypores share similar preferences to the conditions of rotten wood they colonize. At the same time beetle imagines visit sporulating basidiocarps of both the predecessor and successor polypores, and hence spores of successors are present on the body of beetle female actively searching for an appropriate log to lay eggs.

Ostoma ferruginea is a beetle whose larvae live on wood brown-rotted by *Fomitopsis pinicola* and *F. rosea*, as well as on mycelia of certain successor polypores. Both imagines and larvae are often located in the transition between wood and fungus fruit body, and imagines feed on the hymenophore of *Fomitopsis* but also on secondary fungal species, which may colonize logs primarily decayed by *Fomitopsis*, e.g. *Pycnoporellus fulgens* and *Antrodia albobrunnea*. Basidiocarps of *Pycnoporellus fulgens* are often eaten by insect larvae fairly quickly after their development and beetles involved in the dispersal were expected to be found (J. Siitonen, pers. comm.).

Ostoma ferruginea may further be involved in the spore dispersal of *Amylocystis lapponica* often growing together with *Fomitopsis rosea* on the same log. Imagines of *O. ferruginea* visit both of these polypore species. *A. lapponica* and *F. rosea* are known as co-existing species although no predecessor-successor relations were found on the basis of basidiocarp records. *A. lapponica* fruit bodies never arise from the dead basidiocarps of *F. rosea*, and their parallel occurrence may be based on similar substrate preferences.

We believe that at least in some cases spores of successor polypores are dispersed both by wind and beetles, improving chances of rare successor polypores to maintain viable populations.

4.6 Calculations

Fresh basidiocarps attract the highest number of beetle species, collected as imagines (decomposition stage I, Figs. 5, 11). The decline in the number of species (imagines) during the decomposition of the fruit body is interrupted in stage III, when anamorphic fungi start to grow over the fruit body and attract specialized visitors, but structural changes and the decline of basidiocarp

volume are not yet drastic. At the same time stages I and III are favoured by the larvae of fungivore beetles: *Hallomenus* and *Abdera* generally in stage I, while *Dorcatoma* and *Ennearthron* in stage III. These two peaks show that larvae of polypore-dwelling beetles use a strategy either of a short life cycle starting in living fruit body, or a more long one in recently dead polypore basidiocarp. *Cis* larvae and imagines were found in all the stages (Figs. 5–8, 11).

4.7 Notes on selected beetles

Of the two Leiodidae species found, only *Agathidium pisanum* was observed feeding on the hymenophore surface of the polypore *Rigidoporus corticola*. *A. arcticum* imagines were found on *Reticularia* sp., a slime mould.

Imagines of *Phloeocharis subtilissima* hide inside the tubes or wander on the surface of the hymenophore of *Phellinus pini*. Vibration of substrate makes beetles escape in the tubes, where they possibly consume spores. Similar behaviour was shown for another staphylinoid (Ptiliidae) beetle *Baranowskiella ehnstromi* Sörensson, 1997 in the much more fine tubes of *Phellinus conchatus* (Sörensson 1997).




Fig. 11. Numbers of beetle species during the decomposition of fungal basidiocarps. The study was made with polypores of the North Karelian Biosphere Reserve, Finland. For the decomposition stages (I–IV) see Table 1.

Imagines of *Atheta boleticola* were found on the hymenophore of *Phaeolus schweinitzii* together with numerous unidentified staphylinoid larvae living inside the spongy context. Rearings failed, but it is possible that larvae of *A. boleticola* feed on this polypore. Most of Staphylinidae were found on the hymenophore of various polypores, usually moistened by wet soil, dew, or atmospheric precipitation. However, the larval host preferences of the Staphylinidae would deserve further study.

Two related *Cis* species are known to develop in basidiocarp context and on the transition layer between hymenophore and context (Schigel 2002), which was supported in our study with other substrates: *Cis comptus* in *Amylocystis lapponica* and *Gloeophyllum sepiarium*, and *Cis hispidus* in *Antrodia pulvinascens*, *Ceriporiopsis resinascens* and *Lenzites betulinus*. In *Trametes* it was found that species composition of the Ciidae includes three size classes of beetle larvae (and emerging adults) interlacing their burrows of different diameters simultaneously, as it was found in European Russia (Schigel 2002). These three classes in the North Karelian Nature Reserve are represented by *Cis boleti* (largest burrows), *C. hispidus* (medium-sized) and *Octotenus glabriculus* (narrowest), respectively. Burrows of the larvae merge in the late stages of the fruit body decomposition. Fruit body size limits the spatial distribution of larvae inside the basidiocarp: large larvae of *Cis boleti* are located in fruit bodies of *Trametes ochracea*, *T. pubescens* and *T. velutina* at their thick base (umbo), while larvae of *Cis hispidus* and, in particular, *Octotenus glabriculus* tend to graze inside the inner parts of the basidiocarp margin.

Imagines and larvae of *Cis dentatus* burrow the resupinate but fairly thick fruit bodies of *Sarcoporia salmonicolor* and aggregate in their dryer and thicker parts. *C. punctulatus* larvae develop usually in *Trichaptum fuscoviolaceum* and *T. abietinum* (Kompantsev 1982, Schigel 2002) but we reared a few individuals also from *T. pargamenum*, which is typically colonized by other ciids.

Dolichocis laricinus larvae develop in the strongly decomposed fruit bodies of *Antrodia pulvinascens* and *Polyporus leptoccephalus* (*P. varius*).

Imagines of *Rhizophagus dispar*, *Acrulia inflata* and *Orthoperus atomus* were found on the surface of *Phellinus laevigatus* around excrements of unidentified Tineidae larvae. Imagines of *R. dispar* on the fruit bodies of *Gelatoporia*

pannocincta were usually found on excrements of Diptera and Lepidoptera larvae. Processed fungal substrate is more attractive to certain beetle generalists than the hard hyphae of an intact hymenophore. In particular imagines of *Rhizophagus* tend to occur on sporulating polypores (*Inonotus obliquus* and *Fomitopsis pinicola*) but decompose also other fungal substrates, especially euagarics (Moncalvo et al. 2002) such as *Pleurotus* species and *Hypsizygus ulmarius*. *Rhizophagus* was found both on basidiocarps on standing trees and on fruit bodies fallen on the ground. Generally, unlike polypores, euagarics attract less beetles from families other than Staphylinidae, although in Europe the beetle fauna of these two large groups of fungi partly overlap.

Colonization of the fruit bodies of *Phellinus* (*P. pini*, *P. populicola*, *P. igniarius* complex) usually starts when the area of living hymenophore starts to shrink. From this moment the upper part of a fruit body begins to die off. This stage of decomposition (III) is preferred by *Dorcatoma* larvae, for instance *Dorcatoma dresdensis* in *Phellinus lundellii*. Fruit bodies of *P. laevigatus* are sometimes quite thin, less than 1 cm, and in this case the sickle-shaped larvae of *Dorcatoma* change their typical vertical position to horizontal, but still avoid the hymenophore. In dead fruit bodies larvae can consume also the hymenophore. During the whole development cycle the larvae of *Dorcatoma* tend to avoid the outer 1-cm zone of the fruit body. This avoided distance is greater still in the fruit bodies of *P. populicola* because of the deep cracks of the upper side of the fungus. On the one hand, avoiding outer layers of the substrate is characteristic for nearly all the substrate dwellers because of the lower risk of parasite attack and more stable substrate conditions deeper in, but on the other hand imagines emerging from pupae and leaving the fungus face certain difficulties to come out. However, Anobiidae are adapted to live in hard and dry substrates, and Dorcatominae occupy the niche of decomposers of robust polypores. They can also appear at the early stages of fruit body decomposition, thus avoiding competition with Ciidae and other Coleoptera.

Imagines and larvae of *Ennearthron cornutum* are located between the living hymenophore of *Phellinus pini* and the half-dead context of this long-persisting fungus, or in senescent inner layers of the hymenophore. Similar distribu-

tion was found in *P. conchatus* and *P. chrysoloma*, where larvae are using mostly the marginal parts of the fruit body. *P. viticola* is softer in its consistency, and harbours larvae close to the context. On the contrary, *Dichomitus squalens* is a soft polypore, and *Ennearthron cornutum* larvae are restricted to the context. *E. cornutum* is a generalist beetle, but the preferred parts of fruit bodies vary according to fungus species.

The more firm and structured the polypore fruit body is, the more complex the structure of spatial distribution of larvae inside the fungal substrate tends to be. Soft and homogeneous context will lead in an even distribution of larvae: *Ennearthron cornutum* in *Dichomitus squalens*; larvae of *Sulcacis affinis* in the context of *Pyconoporus cinnabarinus*; *Hallomenus* sp. larvae in *Postia alni*, *P. fragilis*, *P. lateritia*, *P. leucomallella*, *P. placenta*, *P. stiptica*, *P. tephroleuca* and *Amylocystis lapponica*.

Dromius sigma, a small, common carabid species, is an inhabitant of lake- and riversides, living under leaves of *Alnus* and *Salix*. Sometimes occasional individuals move far from water in untypical habitats, like the polypore *Cerrena unicolor*.

Acknowledgements: Parts of the Coleoptera material were identified by Dr. N.B. Nikitsky (Zoological Museum of M.V. Lomonosov Moscow State University, Russia), Mr. A.V. Kompantsev and Mrs. T.V. Kompantseva (A.N. Severtsov Institute of Ecology and Evolution, Moscow, Russia), and Mr. V.B. Semenov (E. I. Marcinovsky Institute of Medical Parasitology and Tropical Medicine). We are grateful for this assistance. We appreciate the valuable comments of Dr. Hans Silfverberg (Finnish Museum of Natural History). The Finnish Forest and Park Service kindly invited us to carry out this study and provided facilities for the field work. E-mail discussions in the Finnish coleopterist mailing list "Kuoiraiposti" helped to solve a problem in beetle synonymy. A research grant from the Ministry of Environment, Finland (YM131/5512/2002) is gratefully acknowledged.

References

- Ahti, T., Hämet-Ahti, L. & Jalas, J. 1968: Vegetation zones and their sections in northwestern Europe. – *Ann. Bot. Fennici* 5:169–211.
- Andersen, J., Olberg, S. & Haugen, L. 2000: Saproxyllic beetles (Coleoptera) of Troms and Western Finnmark, northern Norway with exceptional distribution in Fennoscandia. – *Norwegian J. Entomol.* 47: 29–40.
- Benick, L. 1952: Pilzkäfer und Käferpilze. Ökologische und statistische Untersuchungen. – *Acta Zool. Fennica* 70: 1–309.
- Bondarceva, M.A., Krutov, V.I., Lositskaya, V.M. & Svishch, L.G. 1995: Aphyllophoraceous fungi in some pine forests in Eastern Finland and Russian Karelia. – In: Hokkanen, T. J. & Leshko, I. (eds.), *Karelian Biosphere Reserve Studies*. North Karelian Biosphere Reserve, Joensuu: 195–201.
- Bondartseva, M.A., Lositskaya, V.M. & Hokkanen, T.J. 1999: Aphyllophoroid fungi of the North Karelian Biosphere Reserve (Finland). – *Kew Bull.* 54: 589–603.
- Bondartseva, M., Lositskaya, V., Hokkanen, T. & Krutov, V. 2001: Studies on Aphyllophoraceous fungi in North Karelian Biosphere Reserve 1993–1996. – *Nature Protection Publ. Finnish Forest and Park Service A* 131: 91–111.
- Bondartseva, M.A., Lositskaya, V.M. & Zmitrovich, I.V. 1998: Aphyllophoroid fungi of old and primeval forests in the Kotavaara site of North Karelian Biosphere Reserve. – *Folia Cryptog. Estonica* 33: 19–24.
- Bondartseva, M.A., Lositskaya, V.M. & Zmitrovich, I.V. 2000: *Punctularia strigosozonata* (Punctulariaceae) in Europe. – *Karstenia* 40: 9–10.
- Donisthorpe, H. 1931: Coleoptera found in the "birch bracket" fungus *Polyporus betulinus*. – *Entomol. Res.* 43: 14.
- Donisthorpe, H. 1935: The British fungicolous Coleoptera. – *Entomol. Monthly Mag.* 71: 21–31.
- Ehnström, B. & Axelsson, R. 2002: Insektsnag i bark och ved. – *ArtDatabanken SLU*: 165–498.
- Hansen, L. & Knudsen, H. 1997: Nordic Macromycetes 3. Heterobasidioid, aphyllophoroid and gastromycetoid Basidiomycetes. – *Nordsvamp*, Copenhagen. 444 pp.
- Hokkanen, T.J. & Ieshko, E. 1995: Karelian Biosphere Reserve studies. – *North Karelian Biosphere Reserve*, Joensuu. 278 pp.
- Jonsell, M. & Norlander, G. 1995: Field attraction of Coleoptera to odours of the wood-decaying polypores *Fomitopsis pinicola* and *Fomes fomentarius*. – *Ann. Zool. Fennici* 32: 391–402.
- Jonsell, M. & Nordlander, G. 2002: Insects in polypore fungi as indicator species: a comparison between forest sites differing in amounts and continuity of dead wood. – *Forest Ecol. Managem.* 157: 101–118.
- Köljalg, U. & Renvall, P. 2000: *Hydnellum gracilipes*, a link between stipitate and resupinate Hymenomycetes. – *Karstenia* 40:71–77.
- Komonen, A., Siitonen, J. & Mutanen, M. 2001: Insects inhabiting two old growth forest polypore species. – *Entomol. Fennica* 12: 1–14.
- Kompantsev, A.V. 1982: (Morpho-ecological peculiarities of Erotylidae (Coleoptera) larvae – inhabitants of the fruit bodies of higher fungi). – In: *Morpho-ecological adaptations of insects in terrestrial communities*: 81–91. Nauka, Moskva. (In Russian.)
- Kotiranta, H. & Niemelä, T. 1996: Uhanalaiset käävät Suomessa. 2nd revised edition. – *Edita*, Helsinki. 184 pp.
- Lawrence, J. F. 1973: Host preference in Ciid beetles (Coleoptera: Ciidae) inhabiting the fruiting bodies of Basidiomycetes in North America. – *Bull. Mus. Comparative Zool.* 145: 163–212.
- Martikainen, P. 2001: Conservation of threatened saproxyllic beetles: significance of retained aspen *Populus tremula* on clearcut areas. – *Ecol. Bull.* 49: 205–218.

- Moncalvo, J.M., Vilgalys, R., Redhead, S.A., Johnson, J.E., James, T.Y., Aime, M.C., Hofstetter, V., Verduin, S.J.W., Larsson, E., Baroni, T.J., Thorn, R.G., Jacobsson, S., Cléménçon, H. & Miller, O.K. Jr. 2002: One hundred and seventeen clades of euagarics. – *Molecular Phylogen. Evolution* 23: 357–400.
- Müller, C., Jaeger, B., Kompantsev, A.V. & Uhlig, M. 2001: Type and species catalogue of the minute tree-fungus beetles of the Museum für Naturkunde in Berlin, with general information on the Coleoptera collection, its curation and “Historical collection” (Coleoptera, Polyphaga, Ciidae and Pterogeniidae). – *Mitt. Mus. Naturkund. Berlin, Zool. Reihe* 77: 303–323.
- Niemelä, T. 1980: On Fennoscandian polypores 7. The genus *Pycnoporellus*. – *Karstenia* 20: 1–15.
- Niemelä, T. 2003a: What is the commonest polypore in Finland? – *Sieniilehti* 55: 46–53. (In Finnish.)
- Niemelä, T. 2003b: Collecting *Punctularia strigosozonata* (Corticaceae) in Finland and Estonia. – *Sieniilehti* 55: 67–70. (In Finnish.)
- Niemelä, T. 2004: Guide to the polypores of Finland. 15th revised edition. – *Bot. Bull. Univ. Helsinki* 184: 1–148.
- Niemelä, T. & Dai, Y.C. 1998: Luoston suojelualueen kääpäinventointi 1998. – *Metsähallitus, Perä-Pohjolan luontopalvelut, Helsinki*. 32 pp.
- Niemelä, T. & Dai, Y.C. 1999: Polypores of Luosto. – *Nature Protection Publ. Finnish Forest Park Service A 105*: 1–59.
- Niemelä, T. & Kinnunen, J. 2001: Korouoman suojelualueen kääpäinventointi 2001. – *Metsähallitus, Perä-Pohjolan luontopalvelut, Helsinki*. 38 pp.
- Niemelä, T. & Kinnunen, J. 2002: Ylläksen-Aakenuksen suojelualueen kääpäinventointi 2001. – *Metsähallitus, Perä-Pohjolan luontopalvelut, Helsinki*. 30 pp.
- Niemelä, T. & Kinnunen, J. 2003: Pisavaaran luonnonpuiston kääpäinventointi 2003. – *Metsähallitus, Perä-Pohjolan luontopalvelut, Helsinki*. 16 pp.
- Niemelä, T., Kinnunen, J. & Manninen, O. 2002: Koitajoen Natura-alueen kääpäinventointi 2002. – *Metsähallitus, Itä-Suomen luontopalvelut, Helsinki*. 41 pp.
- Niemelä, T., Kinnunen, J., Renvall, P. & Schigel, D. 2003: *Phellodon secretus* (Basidiomycota), a new hyd-naceous fungus from northern pine woodlands. – *Karstenia* 43: 37–44.
- Niemelä, T., Renvall P., Penttilä R. 1995: Interactions of fungi at late stages of wood decomposition. – *Ann. Bot. Fennici* 32: 141–152.
- Nikitsky, N.B. & Schigel, D.S. 2004: Beetles in polypores of the Moscow region: checklist and ecological notes. – *Entomol. Fennica* 15: 6–22.
- Økland, B. 1995: Insect fauna compared between six polypore species in a southern Norwegian spruce forest. – *Fauna Norvegica B* 42: 21–46.
- Palm, T. 1951: Die Holz- und Rindenkäfer der nord-schwedische Laubbäume. – *Meddel. Statens Skogsforskningsinst.* 40: 1–242.
- Palm, T. 1959: Die Holz- und Rindenkäfer der süd- und mittelschwedischen Laubbäume. – *Opusc. Entomologica Suppl.* 16: 1–374.
- Pavior-Smith, K. 1960: The fruiting bodies of macrofungi as habitats for beetles of the family Ciidae (Coleoptera). – *Oikos* 11: 1–17.
- Pouzar, Z. 1971: Notes on taxonomy and nomenclature of *Ischnoderma resinomum* (Fr.) P. Karst. and *I. benzoinum* (Wahlenb.) P. Karst. (Polyporaceae). – *Česká Mykologie* 25: 15–21.
- Rassi, P., Alanen, A., Kanerva, T. & Mannerkoski, I. (eds.) 2001: The 2000 Red List of Finnish species. – *Ministry of the Environment & Finnish Environment Institute*: 1–432.
- Saalas, U. 1917: Die Fichtenkäfer Finnlands. – *Ann. Acad. Sci. Fenniae A* 8: 1–547.
- Saalas, U. 1923: Die Fichtenkäfer Finnlands. – *Ann. Acad. Sci. Fenniae B* 22: 1–746.
- Schigel, D.S. 2002: (Beetle complexes in polypore fungi in East European Plain and Crimea). – *Bull. Moscow Soc. Naturalists* 107: 8–21. (In Russian.)
- Shubin, V.I. & Krutov, V.I. 1979: Griby Karelii i Murmanskoy oblasti. – *Nauka, Leningrad*. 104 pp.
- Siitonen, J., Penttilä, R. & Kotiranta, H. 2001: Coarse woody debris, polyporous fungi and saproxylic insects in an old-growth forest in Voldozero National Park, Russian Karelia. – *Ecological Bulletins* 49: 231–242.
- Silfverberg, H. 1992: Enumeratio Coleopterorum Fennoscandiae, Daniae et Baltiae. – *Helsingin Hyönteisvaihtoyhdystys, Helsinki*. 94 pp.
- Sörensson, M. 1997: Morphological and taxonomical novelties in the world’s smallest beetles, and the first Old World record of *Nanosellini* (Coleoptera: Ptiliidae). – *Systematic Entomol.* 22: 257–283.
- Thunes, K.H. 1994: The coleopteran fauna of *Piptoporus betulinus* and *Fomes fomentarius* (Aphylllophorales: Polyporaceae) in western Norway. – *Entomol. Fennica* 5: 157–168.
- Thunes, K.H., Midtgaard, F. & Gjerde, I. 2000: Diversity of coleoptera of the bracket fungus *Fomitopsis pini-cola* in a Norwegian spruce forest. – *Biodiversity and Conservation* 9: 833–852.
- Yakovlev, E. B., Nikitsky, N. B. & Sherbakov, A. 2001: Saproxylic Coleoptera of unmanaged mature forests in Koitajoki area. – *Diversity studies in Koitajoki Area (North Karelian Biosphere Reserve, Ilomantsi, Finland). Nature Protection Publ. Finnish Forest Park Service A 131*: 32–71.