

Maculineaalcon limitanea nov. ssp. / Lepidoptera:  
Lycaenidae / from Transsylvania, Rumania

Zsolt Bálint

Abstract: A new subspecies of Maculineaalcon D. & SCH. 1775 described. Some remarks on its habitat, ethology and taxonomic problems are given.

Introduction

There is a great number of Maculineaalcon D. & SCH. from the Eastern Carpathians in the Lepidoptera Collection of the Hungarian Natural History Museum. These specimens are different from the other Carpathian Maculineaalcon D. & SCH. specimens in the form of their wings, ground colour and pattern as well as fenology and ecology.

I read through the referring literature and data and examined the Lycaena material of the Hungarian Natural History Museum and I am convinced that the Maculineaalcon populations of the Eastern Carpathians represent a new undescribed subspecies.

Material

Holotype, ♂ Persányi-hegység, Vargyasi-szoros; 1980. VII. 4., leg. Kovács.

Allotype, ♀ Persányi-hegység, Vargyasi-szoros; 1980. VII. 5., leg. Kovács.

Paratypes, 23 ♂ and 14 ♀ Transsylvania, Gyilkostó; E. VII. 1941., coll. V. Bartha / 1 ♀; Mikóújfalu, Nagypatak; 1975. VI. 29., leg. Kovács / 2 ♂, 1 ♀; Uzonkafürdő; 1975. VII. 18., leg. Kovács / 1 ♂; Mikóújfalu, Nagypatak; 650 m; 1976. VII. 1., leg. Kovács / 1 ♂; Békási-szoros, 1100 m; 1978. VII. 16., leg. Kovács / 1 ♀; Békási-szoros, 1000 m; 1979. VI. 25., leg. Kovács / 2 ♂; Vargyasi-szoros, 700 m; 1980. VII. leg. Kovács / 1 ♂, 2 ♀; Vargyasi-szoros, 700 m; 1980. VII. 5., leg. Kovács / 2 ♂, ♀; Vargyasi-szoros, 700 m; 1980. VII. 6., leg.

Kovács /5 ♂, 2 ♀/ Erdély: Békás-szoros, Kupás-patak; 1980.  
VII. 21., leg. Bálint and Ronkay G. /5 ♂, 2 ♀/ Gyergyó, Csi-  
szér; 1980. VIII. 10., leg. Kovács /1 ♀/ Gyergyó, Borzont;  
1980. VII. 13., leg. Kovács /2 ♂, 1 ♀/; Csíki-havasok, Poty-  
tyond; 1981. VII. 19., leg. Bálint /2 ♂/; Románia, Gyilkostó;  
1981. VII. 24-26., leg. Bálint /1 ♂, 1 ♀/; in coll. Hung. Nat.  
Hist. Mus. Budapest /Hungary/

### Description

Males: Alar expanse of Holotype 31 mm. Variation on type series 25 - 35 mm. Wings long and narrow. Upper side of wings deep blue with a delicate black suffusing. Wide black marginal border flowing into the post discal area near the veins but never separating sharply from the ground colour. Most of the specimens has a discoidal spot.



Underside's ground colour dark, brownish grey. Outer line of submarginal spots indistinct, inner line well marked, brownish black. Postdiscal spots small, rings grey and not well defined. Spot in discal cell of fore wing sometimes missing, sometimes only on one side. Fringes white.

/ The upperside of some specimens have a grey ground colour without brownish shade. The postdiscal spots are bigger, rings are whiter and can be better seen on these specimens. /



Females: Alar expanse of Allotype 33 mm. Variation in type specimens 28 - 38 mm. Wings as the male's, but rounder. Upperside of wings greyish brown, totally ground coloured / rarely there are some specimens which has a few blue scales on the base of underwings /. Postdiscal spot line shining softly through.

Underside ground colour greyish brown with rufous shade. Submarginal pattern well marked. Postdiscal spots of fore wings smaller and rings darker as on hind wing. On the base single silvery green scales. Fringes grey.



/ There is only one female specimen on the type series, whose basal area of upperside is blue /Erdély: Békás-szoros, Kupás-patak; 1980. VII. 21., leg. Bálint and Ronkay G./ Its wing form





1. *Maculinea alcon alcon* D. & SCH. 1775 ♂<sup>1</sup>  
Budapest, Rómaifürdő; 1951 VII. 15., leg. dr. Szabó  
underside



3. *Maculinea alcon alcon* D. & SCH. 1775 ♀  
same locality data
4. underside




5. *Maculinea alcon limitanea* ssp. HOLOTYPE ♂  
6. underside



7. *Maculinea alcon limitanea* ssp. ALLJOTYPE ♀

8. underside





- A) male genitalia of *Maculinea alcon alcon* D. & SCH.  
Budapest, Rómaifürdő; gen. prep. Bálint, NO.
- B) male genitalia of *Maculinea alcon limitanea* ssp.  
Transsylvania, Vargyasi-szoros; gen. prep. Bálint, NO. 17.
- C) left valva of *Maculinea alcon alcon* and *Maculinea alcon limitanea*
- D) right valva of *Maculinea alcon alcon* and *Maculinea alcon limitanea*

and underside is similar to the others. This variation is very  
/

Male genitalia: Like alcon alcon. Its size similar, but valve narrower and apex not so curved. Terminal tooth of valve wider and shorter, terminal incision bigger.


#### Penologic and ecologic notes

The habitats of limitanea nov. ssp. can be divided into three groups:

- 1/ subalpine xerophilous meadows on chalk and limestones  
/rocky grass-slopes, talus slopes/
- 2/ montanic mesophilous meadows on vulcanic stones  
/meadow-lands wood clearings/
- 3/ hygrophilous meadows  
/bogs, only two places in Gyergyó /Gheorghieni/ Basin,  
Borzont and in Háromszéki Basin, Rétyi Nyír/ judetul Covasna

We met limitanea mostly on rocky grass-slopes and talus slopes /Persányi-hegység/Muntii Persani/, Békás-szoros /Cheile Bicazului/, where its flying period lasts from the first week of July to the beginning of August. The adults can be found in the surroundings of the foodplant Gentiana cruciata L. The butterfly fauna is rich: Parnassius apollo transsylvanicus SCHWEITZER 1912, Melitaea didyma ESP. 1779, Maculinea arion L. 1758, Plebicula dorylas magna BÁLINT 1985, Pyrgus alveus BHN. 1803 also flies here. On the mesophilous meadows /Csiszér, near Gheorghieni/ the new subspecies is living in a different butterfly association. Its interesting members in the aspect of July are the following: Clossiana titania transsylvanica TILTSCHER, 1913, Brenthis ino ROTT. 1775, Erebia ligea carthusianorum FRUHST. 1909, Palaeochrysophanus hippothoë eurydice ROTT. 1775, Plebicula dorylas magna BÁLINT 1985. Hardly anybody collected in its hygrophilous habitat where the specimens of limitanea are smaller. It is worthy of note that in the birch fen of Háromszéki Basin /judetul Covasna: Rétyi Nyír/ it flies together with Maculinea teleius BERGSTR. 1779 and Colias myrmidone ESP. 1781. This district is warmer as the Gyergyó /Gheorghieni/ Basin which is





9. 10. Rocky grass-slopes /Békás-szoros /Cheile Bicazului,': Kupás-patak/, habitat of *Parnassius apollo transsylvanicus*, *Maculineaalcon limitanea*, ssp. *Maculinea arion*, *Plebicula dorylas*

the coolest place of the Carpathian Basin. The females of alcon limitanea lay eggs on Gentiana pneumonanthe L. as it was observed by Sándor Kovács and Zoltán Kovács in Retyi Nyír.


#### Nomenclatoric and taxonomic notes

Hormuzaki /1897/ mentions that Maculinea alcon is wide-spread and frequent in the hills of Bukowina, above 400 m. He lists many habitats from the Transcarpathians and describes a new aberration with the name ab. cecinae. He thought that the alcon populations of Bukowina belongs to the nominate race. He gives a description of a male alcon from the Rárau Mountains and he writes that this specimen may be a new "geographical race". The descriptions of the specimens from Bukowina and Rárau Mountains are not clear, so we can not know how far the habitat of limitanea ranges. The new subspecies resembles to ssp. sevastos RBL. & Z. 1934 which is described from Macedonia. But the wings of sevastos are wider, the ground colour of males is violescent blue and that of females warm brown. The markings of the under-side of sevastos is clearer, the rings and the spots are more distinct. Ssp. tolistus FRUHST. 1917 from Bosnia has a blue base on the females. Its description is similar to that of xerophila BERGER 1946 and curiosa SZABÒ 1956. I do not share the opinion of Varga /1961/ and Fazekas /1984/. They mention tolistus from Transsylvania and South Transdanubia. It does not occur in the Carpathian Basin.

Maculinea alcon still flies in many places in Transsylvania / Bálint 1980; Coichia & Barbu 1982; Czekelius 1917; Diószeghy 1929/30; Haltrich 1982; Kovács, S. & Kóvacs, 1982; Kónig 1975 and 1982; Popescu-Gorj 1964; Rotschild 1912; /, but it's impossible to determinate their subspecific status because we do not have enough material.

#### Acknowledgements

I wish to express my gratitude to my wife Annamária Kertész for correcting the English manuscript, to Sándor Kovács and Zoltán Kovács / Sfintu Gheorghe, Rumania/ for presenting their material



Distribution map of *Maculineaalcon limitanea*  
Transsylvania, Rumania


- |                                |                             |
|--------------------------------|-----------------------------|
| A) Gyergyótölgyes, Holló-patak | Tulghes, Valea Corbu        |
| B) Gyilkostó, Békásszoros      | Lacu Rosu, Cheile Bicazului |
| C) Gyergyó, Borzont            | Gheorghieni, Borzont        |
| D) Gyergyó, Csiszér            | Gheorghieni, Csiszér        |
| E) Székelyudvarhely            | Odorheiul Secuiesc          |
| F) Csíkménaság, Pottyond       | Armaseni, Potiond           |
| G) Tusnád                      | Tusnad                      |
| H) Baróti-hegység              | Muntii Baraolt              |
| I) Persányi-hegység:           | Muntii Persani:             |
| Vargyasi-szoros                | Cheile Virghisului          |
| J) Bodoki-hegység              | Muntii Bodoc                |

and for their kind help, and also to László Peregovits for taking the photos.

### References

- Bálint, Zs. /1980/: Adatok a nagylepkek elterjedéséhez Erdélyben  
L. /Lepidoptera/ Folia ent. hung. XLI. p: 363 366.
- Berger, L. /1946/: Maculinea Rebeli Hirschke, bona species  
Lambillionea, XLVI. 6-10., p: 95 110
- Coichia, V. & Barbu, A. /1980/: Catalogul colectiei de Lepidoptere  
"N. Delvig" a Muzeului Judetean Brasov Cumidava XII/4,  
Stiintele naturii Brasov, p: 93.
- Czekelius, D. /1917/: Beiträge zur Schmetterlingsfauna Siebenbürgens  
Verhandl. und Mitteil. des Siebenbürgischen Vereins für  
Naturwissenschaften zu Hermannstadt, LXVII., p: 17.
- Diószeghy, L. /1929/1930/: Die Lepidopterenfauna des Retezat-  
Gebirges Verhandl. und Mitteil. des Siebenbürgischen Ver-  
eins für Naturwissenschaften zu Hermannstadt, LXXXIX/LXXX.,  
p: 31.
- Fazekas, I. /1984/: A Keleti-Mecsek lepkefaunája VI., Sikonda  
Zygaenidae, HesperIIDae és Papilionidae faunája /Lepidoptera/  
Folia Comboensis Tom.I., p: 107.
- Fruhstorfer, H. /1917/: Neue palaarktische Lycaeniden Iris,  
XXXI., p: 24 25.
- Haltrich, A. /1982/: Beiträge zur Kenntnis der Rhopaloceren des  
Sălard-Tales /Gurghiu-Gebirge, Ostkarpaten / Studii si  
comunicări II., p: 356.
- Hormuzaki, V. C. /1897/: Die Schmetterlinge /Lepidoptera/ der  
Bukowina Zool. bot. Gesellschaft, XLVII., p: 137.
- Kovács, S. & Kovács, Z. /1982/: Lepidoptere diurne din jurul  
orasului Sfântu Gheorghe Studii si comunicări II., p: 264  
265.
- König, F. /1975/: Catalogul colectiei de lepidoptere a Muzeului  
Banatului Timisoara, p: 214.
- König, F. /1982/: Montane, subalpine, alpine und boreoalpine  
Schmetterlings-Arten aus den Rumänischen Karpaten Studii si  
comunicări II., p: 231.

- Popescu-Gorj, A. /1964/: Catalogue de la collection de lépidoptères "Prof. A. Ostrogovich" du Muséum d'Histoire Naturelle "Grigore Antipa" Bucarest Bucuresti, p: 237.
- Rebel, H. & Zerny, H. /1934/: Wissenschaftliche Ergebnisse der im Jahre 1918 entsendeten Expedition nach Nordalbanien. Die Lepidopterenfauna Albaniens /mit Berücksichtigung der Nachbargebiete/ Akademie der Wissenschaften in Wien, Denkschriften 103., p: 82.
- Rotschild, N.Ch. /1912/: Beitrag zur Lepidopterenfauna der Mezőség Verhandl. und Mitteil. des Siebenbürgischen Vereins für Naturwissenschaften zu Hermannstadt, LXII., p: 7.
- Szabó, R. /1956/: Magyarország Lycaenidái Folia ent. hung. TOM.IX/series nova/ 13., p: 262 264.
- Varga, Z. /1961/: Allatföldrajzi vizsgálatok az Északborsodi Karszt nagylepke-faunáján Folia ent. hung. TOM. XIV/series nova/. 28., p: 352 356.



11. Vargyasi-szoros /  
Muntii Persani:  
Cheile Virghisului/  
Locus-typicus of  
Maculineaalcon  
Limitanea n.

Address of author: Zsolt Bálint  
Természettudományi Múzeum Allattára  
Budapest VIII  
Baross utca 13.  
H 1088

# ZOBODAT - [www.zobodat.at](http://www.zobodat.at)

Zoologisch-Botanische Datenbank/Zoological-Botanical Database

Digitale Literatur/Digital Literature

Zeitschrift/Journal: [Galathea, Berichte des Kreises Nürnberger Entomologen e.V.](#)

Jahr/Year: 1985

Band/Volume: [1](#)

Autor(en)/Author(s): Balint [Bálint] Zsolt

Artikel/Article: [Maculineaalcon limitanea nov. ssp. / Lepidoptera: Lycaenidae / from Transsylvania, Rumania 62-74](#)